

GENERATION OF RAY CLASS FIELDS MODULO 2, 3, 4 OR 6 BY USING THE WEBER FUNCTION

HO YUN JUNG, JA KYUNG KOO, AND DONG HWA SHIN

ABSTRACT. Let K be an imaginary quadratic field with ring of integers \mathcal{O}_K . Let E be an elliptic curve with complex multiplication by \mathcal{O}_K , and let h_E be the Weber function on E . Let $N \in \{2, 3, 4, 6\}$. We show that h_E alone when evaluated at a certain N -torsion point on E generates the ray class field of K modulo $N\mathcal{O}_K$. This would be a partial answer to the question raised by Hasse and Ramachandra.

1. Introduction

Let K be an imaginary quadratic field with ring of integers \mathcal{O}_K . Corresponding to the elliptic curve E in $\mathbb{P}^2(\mathbb{C})$ with parametrization

$$\begin{aligned} \varphi_E : \mathbb{C}/\mathcal{O}_K &\xrightarrow{\sim} E : y^2 = 4x^3 - g_2(\mathcal{O}_K)x - g_3(\mathcal{O}_K) \\ z &\mapsto [\wp(z; \mathcal{O}_K) : \wp'(z; \mathcal{O}_K) : 1] \end{aligned}$$

(see §2), the *Weber function* $h_E : E \rightarrow \mathbb{P}^1(\mathbb{C})$ is defined by

$$(1) \quad h_E(x, y) = \begin{cases} (g_2g_3/\Delta)x & \text{if } K \neq \mathbb{Q}(\sqrt{-1}), \mathbb{Q}(\sqrt{-3}), \\ (g_2^2/\Delta)x^2 & \text{if } K = \mathbb{Q}(\sqrt{-1}), \\ (g_3/\Delta)x^3 & \text{if } K = \mathbb{Q}(\sqrt{-3}), \end{cases}$$

where $g_2 = g_2(\mathcal{O}_K)$, $g_3 = g_3(\mathcal{O}_K)$ and $\Delta = g_2^3 - 27g_3^2$. This function gives rise to an isomorphism of the quotient variety $E/\text{Aut}(E)$ onto $\mathbb{P}^1(\mathbb{C})$ ([15, Theorem 7 in Chapter 1]).

For a proper nontrivial ideal \mathfrak{m} of \mathcal{O}_K , let $K_{\mathfrak{m}}$ denote the ray class field of K modulo \mathfrak{m} . As a consequence of the theory of complex multiplication, Hasse proved in [5] that

$$K_{\mathfrak{m}} = K(j(E), h_E(P)) \text{ for some } \mathfrak{m}\text{-torsion point } P \text{ on } E,$$

Received March 30, 2017; Revised August 16, 2017; Accepted September 5, 2017.
2010 *Mathematics Subject Classification*. Primary 11R37; Secondary 11G15, 11G16.

Key words and phrases. class field theory, complex multiplication, Weber function.

The first author was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea Government (MISP) (2016R1A5A1008055 and 2017R1C1B2010652). The third author was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea Government (MISP) (2017R1A2B1006578) and by Hankuk University of Foreign Studies Research Fund of 2017.

where $j(E) = 1728g_2^3/\Delta$ is the j -invariant of E (see also [3, §26]). In his letter to Hecke, Hasse further asked whether every finite abelian extension of K can be generated by a singular value of the Weber function without the j -invariant ([4, p. 91] or [17, p. 105]). He called this a “problem that lies rather deep”, and Artin found this question “very interesting”. On the other hand, in 1933 and 1936, Sugawara ([23] and [24]) first gave a partial answer to this question.

Recently, Koo et al. ([12] and [13]) showed by applying Kronecker’s second limit formula that if $\mathfrak{m} = N\mathcal{O}_K$ for a positive integer $N (> 1)$, then $K_{\mathfrak{m}}$ is generated solely by $h_E(\varphi_E(1/N))$ or $h_E(\varphi_E(2/N))$ except for the four cases where $N = 2, 3, 4, 6$. In this paper, we shall completely resolve these exceptional cases (Remark 4.6 (iv), Theorems 5.1, 5.3, 6.1, 6.5, Corollaries 7.1, 8.1, 8.2, 8.3 and Theorem 8.5) without using L -function arguments. To this end we shall express Fricke functions in terms of Siegel functions, and make use of Shimura’s reciprocity law and inequality arguments.

2. Meromorphic modular functions

In this section, we shall introduce basic properties of Fricke functions and Siegel functions.

For a lattice Λ in \mathbb{C} , let E_{Λ} be the elliptic curve in $\mathbb{P}^2(\mathbb{C})$ given by the affine Weierstrass form

$$E_{\Lambda} : y^2 = 4x^3 - g_2(\Lambda)x - g_3(\Lambda),$$

where

$$g_2(\Lambda) = 60 \sum_{\lambda \in \Lambda \setminus \{0\}} \frac{1}{\lambda^4} \quad \text{and} \quad g_3(\Lambda) = 140 \sum_{\lambda \in \Lambda \setminus \{0\}} \frac{1}{\lambda^6}.$$

Then, there is a complex analytic isomorphism $\varphi_{E_{\Lambda}} : \mathbb{C}/\Lambda \rightarrow E_{\Lambda}$ of complex Lie groups defined by

$$\varphi_{E_{\Lambda}}(z + \Lambda) = [\wp(z; \Lambda) : \wp'(z; \Lambda) : 1],$$

where

$$(2) \quad \wp(z; \Lambda) = \frac{1}{z^2} + \sum_{\lambda \in \Lambda \setminus \{0\}} \left\{ \frac{1}{(z - \lambda)^2} - \frac{1}{\lambda^2} \right\} \quad (z \in \mathbb{C})$$

is the Weierstrass \wp -function relative to Λ and $\wp' = d\wp/dz$ ([19, Proposition 3.6(b) in Chapter VI]). Note that for $z_1, z_2 \in \mathbb{C} \setminus \Lambda$

$$\wp(z_1; \Lambda) = \wp(z_2; \Lambda) \iff z_1 \equiv \pm z_2 \pmod{\Lambda}$$

([2, Lemma 10.4]). Furthermore, the j -invariant of the elliptic curve E_{Λ} is given by

$$(3) \quad j(E_{\Lambda}) = 1728 \frac{g_2(\Lambda)^3}{\Delta(\Lambda)},$$

where $\Delta(\Lambda) = g_2(\Lambda)^3 - 27g_3(\Lambda)^2$.

Proposition 2.1. *Given a lattice $\Lambda = [\lambda_1, \lambda_2]$ in \mathbb{C} , the polynomial*

$$4x^3 - g_2(\Lambda)x - g_3(\Lambda)$$

has three distinct roots $\wp(\lambda_1/2; \Lambda)$, $\wp(\lambda_2/2; \Lambda)$ and $\wp((\lambda_1 + \lambda_2)/2; \Lambda)$.

Proof. See [19, Proposition 3.6(a) in Chapter VI]. \square

Remark 2.2. Thus, $\Delta(\Lambda) = g_2(\Lambda)^3 - 27g_3(\Lambda)^2$, as the discriminant of the polynomial $4x^3 - g_2(\Lambda)x - g_3(\Lambda)$, is always nonzero.

The modular group $\mathrm{PSL}_2(\mathbb{Z}) = \mathrm{SL}_2(\mathbb{Z})/\{\pm I_2\}$ acts on the complex upper-half plane $\mathbb{H} = \{\tau \in \mathbb{C} \mid \mathrm{Im}(\tau) > 0\}$ by fractional linear transformations, namely, for each $\gamma = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathrm{SL}_2(\mathbb{Z})$

$$\gamma(\tau) = \frac{a\tau + b}{c\tau + d} \quad (\tau \in \mathbb{H}).$$

For a positive integer N , let $\Gamma(N) = \{\gamma \in \mathrm{SL}_2(\mathbb{Z}) \mid \gamma \equiv I_2 \pmod{N \cdot M_2(\mathbb{Z})}\}$ be the principal congruence subgroup of $\mathrm{SL}_2(\mathbb{Z})$ of level N , which gives rise to the modular curve

$$X(N) = \bar{\Gamma}(N) \backslash \mathbb{H}^*,$$

where $\bar{\Gamma}(N) = \pm\Gamma(N)/\{\pm I_2\}$ and $\mathbb{H}^* = \mathbb{H} \cup \mathbb{Q} \cup \{i\infty\}$ ([18, Chapter 1]). If $\mathbb{C}(X(N))$ denotes the field of meromorphic functions on $X(N)$, then it is a Galois extension of $\mathbb{C}(X(1))$ with

$$\mathrm{Gal}(\mathbb{C}(X(N))/\mathbb{C}(X(1))) \simeq \mathrm{PSL}_2(\mathbb{Z})/\bar{\Gamma}(N) \simeq \mathrm{SL}_2(\mathbb{Z}/N\mathbb{Z})/\{\pm I_2\}$$

([18, Proposition 6.1]). Observe that every function in $\mathbb{C}(X(N))$ has Fourier expansion with respect to $q^{1/N}$, where $q = e^{2\pi i\tau}$ ([18, §2.1]). Let \mathcal{F}_N be the subfield of $\mathbb{C}(X(N))$ consisting of functions whose Fourier coefficients belong to the N th cyclotomic field $\mathbb{Q}(\zeta_N)$, where $\zeta_N = e^{2\pi i/N}$. Then, \mathcal{F}_N is a Galois extension of $\mathcal{F}_1 = \mathbb{Q}(j(\tau))$ with

$$\mathrm{Gal}(\mathcal{F}_N/\mathcal{F}_1) \simeq \mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})/\{\pm I_2\},$$

where $j(\tau) = j([\tau, 1])$ is the elliptic modular function ([18, Theorem 6.6 and Proposition 6.9(1)]). Furthermore, $\mathrm{Gal}(\mathbb{C}(X(N))/\mathbb{C}(X(1)))$ can be considered as a subgroup of $\mathrm{Gal}(\mathcal{F}_N/\mathcal{F}_1)$.

For $\mathbf{v} = \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} \in \mathbb{Q}^2 \setminus \mathbb{Z}^2$, the Fricke function $f_{\mathbf{v}}(\tau)$ is defined by

$$(4) \quad f_{\mathbf{v}}(\tau) = \frac{g_2(\tau)g_3(\tau)}{\Delta(\tau)} \wp(v_1\tau + v_2; [\tau, 1]) \quad (\tau \in \mathbb{H}),$$

where $g_2(\tau) = g_2([\tau, 1])$, $g_3(\tau) = g_3([\tau, 1])$ and $\Delta(\tau) = g_2(\tau)^3 - 27g_3(\tau)^2$. And, the Siegel function $g_{\mathbf{v}}(\tau)$ is given by the infinite product expansion

$$(5) \quad g_{\mathbf{v}}(\tau) = -e^{\pi i v_2(v_1-1)} q^{(1/2)\mathbf{B}_2(v_1)} (1 - q^{v_1} e^{2\pi i v_2}) \prod_{n=1}^{\infty} (1 - q^{n+v_1} e^{2\pi i v_2}) (1 - q^{n-v_1} e^{-2\pi i v_2}) \quad (\tau \in \mathbb{H}),$$

where $\mathbf{B}_2(x) = x^2 - x + 1/6$ is the Bernoulli second polynomial. We then see that $g_{\mathbf{v}}(\tau)$ has neither zeros nor poles on \mathbb{H} .

Proposition 2.3. *Let N be an integer such that $N \geq 2$, and let $\mathbf{v} = \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} \in (1/N)\mathbb{Z}^2 \setminus \mathbb{Z}^2$.*

(i) *If $\mathbf{u} \in \mathbb{Q}^2 \setminus \mathbb{Z}^2$ satisfies $\mathbf{u} \equiv \mathbf{v}$ or $-\mathbf{v} \pmod{\mathbb{Z}^2}$, then*

$$f_{\mathbf{u}}(\tau) = f_{\mathbf{v}}(\tau) \quad \text{and} \quad g_{\mathbf{u}}(\tau)^{12N/\gcd(6, N)} = g_{\mathbf{v}}(\tau)^{12N/\gcd(6, N)}.$$

(ii) *$f_{\mathbf{v}}(\tau)$ and $g_{\mathbf{v}}(\tau)^{12N/\gcd(6, N)}$ belong to \mathcal{F}_N .*

(iii) *If $\gamma \in \mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})/\{\pm I_2\} \simeq \mathrm{Gal}(\mathcal{F}_N/\mathcal{F}_1)$, then*

$$f_{\mathbf{v}}(\tau)^\gamma = f_{\gamma^T \mathbf{v}}(\tau) \quad \text{and} \quad \left(g_{\mathbf{v}}(\tau)^{12N/\gcd(6, N)} \right)^\gamma = g_{\gamma^T \mathbf{v}}(\tau)^{12N/\gcd(6, N)},$$

where γ^T means the transpose of γ .

(iv) *If \mathbf{v} satisfies $\mathbf{v} \equiv \begin{bmatrix} v_1 \\ dv_2 \end{bmatrix}$ or $-\begin{bmatrix} v_1 \\ dv_2 \end{bmatrix} \pmod{\mathbb{Z}^2}$ for all $d \in (\mathbb{Z}/N\mathbb{Z})^*$,*

then $f_{\mathbf{v}}(\tau)$ and $g_{\mathbf{v}}(\tau)^{12N/\gcd(6, N)}$ have rational Fourier coefficients.

Proof. See [18, (6.1.4), (6.1.6), Proposition 6.1 and Theorem 6.6] and [14, Proposition 1.3 in Chapter 2 and Theorems 5.2 and 5.3 in Chapter 3]. \square

For $r \in \mathbb{R}$, let $\langle r \rangle$ be the fractional part of r in the interval $[0, 1)$.

Proposition 2.4. *Let $\mathbf{v} = \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} \in \mathbb{Q}^2 \setminus \mathbb{Z}^2$.*

(i) *We have the q -order formula*

$$\mathrm{ord}_q g_{\mathbf{v}}(\tau) = \frac{1}{2} \mathbf{B}_2(\langle v_1 \rangle).$$

(ii) *If $\mathbf{u} \in \mathbb{Q}^2 \setminus \mathbb{Z}^2$ with $\mathbf{u} \not\equiv \pm \mathbf{v} \pmod{\mathbb{Z}^2}$, then*

$$(f_{\mathbf{u}}(\tau) - f_{\mathbf{v}}(\tau))^6 = \frac{j(\tau)^2(j(\tau) - 1728)^3 g_{\mathbf{u}+\mathbf{v}}(\tau)^6 g_{\mathbf{u}-\mathbf{v}}(\tau)^6}{2^{30} \cdot 3^{24} g_{\mathbf{u}}(\tau)^{12} g_{\mathbf{v}}(\tau)^{12}}.$$

Proof. (i) See [14, p. 31].

(ii) See [14, p. 51]. \square

3. Fricke and Siegel functions of level 2, 3, 4 or 6

We shall develop several lemmas on Fricke and Siegel functions of low level for later use.

For a positive integer N , let

$$\Gamma_1(N) = \left\{ \gamma \in \mathrm{SL}_2(\mathbb{Z}) \mid \gamma \equiv \begin{bmatrix} 1 & * \\ 0 & 1 \end{bmatrix} \pmod{N \cdot M_2(\mathbb{Z})} \right\},$$

and let $X_1(N) = \bar{\Gamma}_1(N) \backslash \mathbb{H}^*$ be the corresponding modular curve with $\bar{\Gamma}_1(N) = \pm \Gamma_1(N) / \{\pm I_2\}$. Let $\mathbb{C}(X_1(N))$ be the field of meromorphic functions on $X_1(N)$, and let $\mathbb{Q}(X_1(N))$ be its subfield consisting of functions with rational Fourier coefficients. Note that $X_1(N)$ has genus zero if and only if $N \in \{1, 2, \dots, 10, 12\}$ ([9, Theorem 1]).

Proposition 3.1. *If $N \in \{2, 3\}$, then we have*

$$\mathbb{Q}(X_1(N)) = \mathbb{Q}\left(g_{\begin{bmatrix} 0 \\ 1/N \end{bmatrix}}(\tau)^{12}\right).$$

Proof. For any $\gamma = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \Gamma_1(N)$, we derive that

$$\begin{aligned} \left(g_{\begin{bmatrix} 0 \\ 1/N \end{bmatrix}}^{12} \circ \gamma\right)(\tau) &= \left(g_{\begin{bmatrix} 0 \\ 1/N \end{bmatrix}}(\tau)^{12}\right)^\gamma \quad \text{by Proposition 2.3(ii)} \\ &= g_{\gamma\tau\begin{bmatrix} 0 \\ 1/N \end{bmatrix}}(\tau)^{12} \quad \text{by Proposition 2.3(iii)} \\ &= g_{\begin{bmatrix} c/N \\ d/N \end{bmatrix}}(\tau)^{12} \\ &= g_{\begin{bmatrix} 0 \\ 1/N \end{bmatrix}}(\tau)^{12} \quad \text{by Proposition 2.3(i) and the fact } \gamma \in \Gamma_1(N). \end{aligned}$$

This shows that $g_{\begin{bmatrix} 0 \\ 1/N \end{bmatrix}}(\tau)^{12}$ belongs to $\mathbb{C}(X_1(N))$. And, $g_{\begin{bmatrix} 0 \\ 1/N \end{bmatrix}}(\tau)^{12}$ has rational Fourier coefficients by Proposition 2.3(i) and (iv).

The modular curve $X_1(N)$ has two inequivalent cusps $i\infty$ and 0 of width 1 and N , respectively ([10, Lemma 3]). We then see by Proposition 2.4(i) that the order of $g_{\begin{bmatrix} 0 \\ 1/N \end{bmatrix}}(\tau)^{12}$ at the cusp $i\infty$ is

$$\text{width of } i\infty \text{ on } X_1(N) \cdot \text{ord}_q g_{\begin{bmatrix} 0 \\ 1/N \end{bmatrix}}(\tau)^{12} = 1 \cdot \left(12 \cdot \frac{1}{2} \mathbf{B}_2(0)\right) = 1.$$

Since $g_{\begin{bmatrix} 0 \\ 1/N \end{bmatrix}}(\tau)^{12}$ has neither zeros nor pole on \mathbb{H} , its order at the cusp 0 should be -1 . This implies that $g_{\begin{bmatrix} 0 \\ 1/N \end{bmatrix}}(\tau)^{12}$ gives rise to an isomorphism

$$\begin{aligned} X_1(N) &\rightarrow \mathbb{P}^1(\mathbb{C}) \\ \tau &\mapsto \left[g_{\begin{bmatrix} 0 \\ 1/N \end{bmatrix}}(\tau)^{12} : 1 \right], \end{aligned}$$

from which it follows that $\mathbb{C}(X_1(N)) = \mathbb{C}\left(g_{\begin{bmatrix} 0 \\ 1/N \end{bmatrix}}(\tau)^{12}\right)$. Since $g_{\begin{bmatrix} 0 \\ 1/N \end{bmatrix}}(\tau)^{12}$ has rational Fourier coefficients, we conclude by [11, Lemma 4.1] that

$$\mathbb{Q}(X_1(N)) = \mathbb{Q}\left(g_{\begin{bmatrix} 0 \\ 1/N \end{bmatrix}}(\tau)^{12}\right). \quad \square$$

Lemma 3.2. *Let $\tau \in \mathbb{H}$ such that both $g_2(\tau)$ and $g_3(\tau)$ are nonzero. Then, the polynomial*

$$2916 \cdot 1728^3 x^3 - 27 \cdot 1728 j(\tau)(j(\tau) - 1728)x - j(\tau)(j(\tau) - 1728)^2$$

has three distinct roots $f_{\begin{bmatrix} 0 \\ 1/2 \end{bmatrix}}(\tau)$, $f_{\begin{bmatrix} 1/2 \\ 0 \end{bmatrix}}(\tau)$ and $f_{\begin{bmatrix} 1/2 \\ 1/2 \end{bmatrix}}(\tau)$.

Proof. By Proposition 2.1 and the definition (4), the polynomial

$$4 \left(\frac{\Delta(\tau)}{g_2(\tau)g_3(\tau)} x \right)^3 - g_2(\tau) \left(\frac{\Delta(\tau)}{g_2(\tau)g_3(\tau)} x \right) - g_3(\tau)$$

has three distinct roots $f_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix} \right]}(\tau)$, $f_{\left[\begin{smallmatrix} 1/2 \\ 0 \end{smallmatrix} \right]}(\tau)$ and $f_{\left[\begin{smallmatrix} 1/2 \\ 1/2 \end{smallmatrix} \right]}(\tau)$. By using the definition (3), one can express the above polynomial as

$$\begin{aligned} & \left(\frac{\Delta(\tau)}{g_2(\tau)g_3(\tau)} \right)^3 \left(4x^3 - \frac{g_2(\tau)^3 g_3(\tau)^2}{\Delta(\tau)^2} x - \frac{g_2(\tau)^3 g_3(\tau)^4}{\Delta(\tau)^3} \right) \\ &= \left(\frac{\Delta(\tau)}{g_2(\tau)g_3(\tau)} \right)^3 \left\{ 4x^3 - \frac{j(\tau)}{1728} \left(\frac{j(\tau)}{27 \cdot 1728} - \frac{1}{27} \right) x - \frac{j(\tau)}{1728} \left(\frac{j(\tau)}{27 \cdot 1728} - \frac{1}{27} \right)^2 \right\} \\ &= \left(\frac{\Delta(\tau)}{g_2(\tau)g_3(\tau)} \right)^3 \frac{1}{27^2 \cdot 1728^3} \\ & \quad \times \left\{ 2916 \cdot 1728^3 x^3 - 27 \cdot 1728 j(\tau) (j(\tau) - 1728) x - j(\tau) (j(\tau) - 1728)^2 \right\}. \end{aligned}$$

This proves the lemma. \square

Remark 3.3. Let $\tau \in \mathbb{H}$. Then we get that

$$\begin{aligned} g_2(\tau) = 0 &\iff j(\tau) = 0 \iff \tau = \gamma(\zeta_3) \text{ for some } \gamma \in \text{PSL}_2(\mathbb{Z}), \\ g_3(\tau) = 0 &\iff j(\tau) = 1728 \iff \tau = \gamma(i) \text{ for some } \gamma \in \text{PSL}_2(\mathbb{Z}) \end{aligned}$$

([2, Theorem 10.9 and p. 219]).

Lemma 3.4. *We have the relation*

$$j(\tau) = \frac{\left(g_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix} \right]}(\tau)^{12} + 16 \right)^3}{g_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix} \right]}(\tau)^{12}}.$$

Proof. See [2, Theorem 12.17]. \square

Remark 3.5. It follows that

$$j(\tau)^2 (j(\tau) - 1728)^3 = \frac{\left(g_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix} \right]}(\tau)^{12} + 16 \right)^6 \left(g_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix} \right]}(\tau)^{12} + 64 \right)^3 \left(g_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix} \right]}(\tau)^{12} - 8 \right)^6}{g_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix} \right]}(\tau)^{60}}.$$

Lemma 3.6. *We establish the expression*

$$f_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix} \right]}(\tau) = -\frac{1}{31104} \frac{g_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix} \right]}(\tau)^{36} + 72g_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix} \right]}(\tau)^{24} + 384g_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix} \right]}(\tau)^{12} - 8192}{g_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix} \right]}(\tau)^{12}}.$$

Proof. For convenience, set

$$f = f_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix} \right]}(\tau), \quad g = g_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix} \right]}(\tau)^{12} \quad \text{and} \quad j = j(\tau).$$

In a similar way to the first part of the proof of Proposition 3.1, one can show that f belongs to $\mathbb{Q}(X_1(2))$. Furthermore, f satisfies

$$2916 \cdot 1728^3 f^3 - 27 \cdot 1728j(j - 1728)f - j(j - 1728)^2 = 0$$

by Lemma 3.2. It then follows from Lemma 3.4 that

$$\begin{aligned} & 2916 \cdot 1728^3 g^3 f^3 - 27 \cdot 1728g(g + 16)^3 \{(g + 16)^3 - 1728g\}f \\ & - (g + 16)^3 \{(g + 16)^3 - 1728g\}^2 = 0, \end{aligned}$$

which can be factored into

$$\begin{aligned} & (31104gf + g^3 + 72g^2 + 384g - 8192) \\ (6) \quad & \times \{483729408g^2 f^2 + (-15552g^4 - 1119744g^3 - 5971968g^2 + 127401984g)f \\ & - g^6 - 72g^5 - 192g^4 + 20992g^3 - 1966080g + 8388608\} = 0. \end{aligned}$$

Since $\mathbb{Q}(X_1(2)) = \mathbb{Q}(g)$ by Proposition 3.1 and f lies in $\mathbb{Q}(X_1(2))$, we get $\deg(f, \mathbb{Q}(g)) = 1$. Note that the second factor in the left side of (6) is irreducible as a quadratic polynomial in f over $\mathbb{Q}(g)$ because its discriminant

$$\begin{aligned} & (-15552g^4 - 1119744g^3 - 5971968g^2 + 127401984g)^2 \\ & - 4(483729408g^2)(-g^6 - 72g^5 - 192g^4 + 20992g^3 - 1966080g + 8388608) \\ & = 2176782336g^8 + 174142586880g^7 + \cdots + 2282521714753536g^3 \end{aligned}$$

is not a square in $\mathbb{Q}(g)$. Thus we must have

$$31104gf + g^3 + 72g^2 + 384g - 8192 = 0.$$

Therefore, we achieve

$$f = -\frac{1}{31104} \frac{g^3 + 72g^2 + 384g - 8192}{g}. \quad \square$$

Lemma 3.7. *We have*

$$f_{\begin{bmatrix} 0 \\ 1/3 \end{bmatrix}}(\tau) + f_{\begin{bmatrix} 1/3 \\ 0 \end{bmatrix}}(\tau) + f_{\begin{bmatrix} 1/3 \\ 1/3 \end{bmatrix}}(\tau) + f_{\begin{bmatrix} 1/3 \\ 2/3 \end{bmatrix}}(\tau) = 0.$$

Proof. Let $T = \{(a, b) \in \mathbb{Z}^2 \mid 0 \leq a, b \leq 2\}$. For each $(a, b) \in T \setminus \{(0, 0)\}$, we attain by the definition (2) that

$$\begin{aligned} & \wp\left(\frac{a}{3}\tau + \frac{b}{3}; [\tau, 1]\right) \\ & = \sum_{(m, n) \in \mathbb{Z}^2} \frac{9}{\{(a - 3m)\tau + (b - 3n)\}^2} - \sum_{(m, n) \in \mathbb{Z}^2 \setminus \{(0, 0)\}} \frac{1}{(m\tau + n)^2} \\ & = \sum_{\substack{(k, \ell) \in \mathbb{Z}^2 \setminus \{(0, 0)\} \\ k \equiv a \pmod{3}, \ell \equiv b \pmod{3}}} \frac{9}{(k\tau + \ell)^2} - \sum_{\substack{(m, n) \in \mathbb{Z}^2 \setminus \{(0, 0)\} \\ m \equiv n \equiv 0 \pmod{3}}} \frac{9}{(m\tau + n)^2}. \end{aligned}$$

We then derive that

$$\begin{aligned}
& \sum_{(a,b) \in T \setminus \{(0,0)\}} \wp \left(\frac{a}{3}\tau + \frac{b}{3}; [\tau, 1] \right) \\
&= \sum_{(a,b) \in T \setminus \{(0,0)\}} \left(\sum_{\substack{(k,\ell) \in \mathbb{Z}^2 \setminus \{(0,0)\} \\ k \equiv a \pmod{3}, \ell \equiv b \pmod{3}}} \frac{9}{(k\tau + \ell)^2} \right) \\
&\quad - 8 \sum_{\substack{(m,n) \in \mathbb{Z}^2 \setminus \{(0,0)\} \\ m \equiv n \equiv 0 \pmod{3}}} \frac{9}{(m\tau + n)^2} \\
&= \sum_{(a,b) \in T} \left(\sum_{\substack{(k,\ell) \in \mathbb{Z}^2 \setminus \{(0,0)\} \\ k \equiv a \pmod{3}, \ell \equiv b \pmod{3}}} \frac{9}{(k\tau + \ell)^2} \right) \\
&\quad - 9 \sum_{\substack{(m,n) \in \mathbb{Z}^2 \setminus \{(0,0)\} \\ m \equiv n \equiv 0 \pmod{3}}} \frac{9}{(m\tau + n)^2} \\
&= \sum_{(k,\ell) \in \mathbb{Z}^2 \setminus \{(0,0)\}} \frac{9}{(k\tau + \ell)^2} - 9 \sum_{(m,n) \in \mathbb{Z}^2 \setminus \{(0,0)\}} \frac{1}{(m\tau + n)^2} \\
&= 0.
\end{aligned}$$

Thus we obtain by Proposition 2.3(i) and the definition (4) that

$$\begin{aligned}
& f_{\begin{bmatrix} 0 \\ 1/3 \end{bmatrix}}(\tau) + f_{\begin{bmatrix} 1/3 \\ 0 \end{bmatrix}}(\tau) + f_{\begin{bmatrix} 1/3 \\ 1/3 \end{bmatrix}}(\tau) + f_{\begin{bmatrix} 1/3 \\ 2/3 \end{bmatrix}}(\tau) \\
&= \frac{1}{2} \sum_{(a,b) \in T \setminus \{(0,0)\}} f_{\begin{bmatrix} a/3 \\ b/3 \end{bmatrix}}(\tau) \\
&= \frac{g_2(\tau)g_3(\tau)}{2\Delta(\tau)} \sum_{(a,b) \in T \setminus \{(0,0)\}} \wp \left(\frac{a}{3}\tau + \frac{b}{3}; [\tau, 1] \right) \\
&= 0. \quad \square
\end{aligned}$$

Lemma 3.8. *We have*

$$f_{\begin{bmatrix} 0 \\ 1/2 \end{bmatrix}}(\tau) = \frac{1}{2} \left(f_{\begin{bmatrix} 0 \\ 1/4 \end{bmatrix}}(\tau) + f_{\begin{bmatrix} 2/4 \\ 1/4 \end{bmatrix}}(\tau) \right).$$

Proof. Let $T = \{(0, 1), (0, 3), (2, 1), (2, 3)\}$. We attain by Proposition 2.3(i) and the definitions (2) and (4) that

$$\frac{1}{2} \left(f_{\begin{bmatrix} 0 \\ 1/4 \end{bmatrix}}(\tau) + f_{\begin{bmatrix} 2/4 \\ 1/4 \end{bmatrix}}(\tau) \right)$$

$$\begin{aligned}
&= \frac{1}{4} \sum_{(a,b) \in T} f_{\left[\begin{smallmatrix} a/4 \\ b/4 \end{smallmatrix} \right]}(\tau) \\
&= \frac{g_2(\tau)g_3(\tau)}{4\Delta(\tau)} \sum_{(a,b) \in T} \wp \left(\frac{a}{4}\tau + \frac{b}{4}; [\tau, 1] \right) \\
&= \frac{g_2(\tau)g_3(\tau)}{4\Delta(\tau)} \sum_{(a,b) \in T} \left(\sum_{(m,n) \in \mathbb{Z}^2} \frac{16}{\{(a-4m)\tau + (b-4n)\}^2} - \sum_{(m,n) \in \mathbb{Z}^2 \setminus \{(0,0)\}} \frac{1}{(m\tau + n)^2} \right) \\
&= \frac{g_2(\tau)g_3(\tau)}{\Delta(\tau)} \left(\sum_{(m,n) \in \mathbb{Z}^2} \frac{4}{\{-2m\tau + (1-2n)\}^2} - \sum_{(m,n) \in \mathbb{Z}^2 \setminus \{(0,0)\}} \frac{1}{(m\tau + n)^2} \right) \\
&= \frac{g_2(\tau)g_3(\tau)}{\Delta(\tau)} \wp \left(\frac{1}{2}; [\tau, 1] \right) \\
&= f_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix} \right]}(\tau). \quad \square
\end{aligned}$$

Lemma 3.9. *We have*

$$f_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix} \right]}(\tau) = \frac{1}{4} \left(f_{\left[\begin{smallmatrix} 0 \\ 1/6 \end{smallmatrix} \right]}(\tau) + f_{\left[\begin{smallmatrix} 2/6 \\ 1/6 \end{smallmatrix} \right]}(\tau) + f_{\left[\begin{smallmatrix} 2/6 \\ 3/6 \end{smallmatrix} \right]}(\tau) + f_{\left[\begin{smallmatrix} 2/6 \\ 5/6 \end{smallmatrix} \right]}(\tau) \right).$$

Proof. Let $T = \{(0, 1), (0, 3), (0, 5), (2, 1), (2, 3), (2, 5), (4, 1), (4, 3), (4, 5)\}$. We derive by Proposition 2.3(i) and the definitions (2) and (4) that

$$\begin{aligned}
&\frac{1}{4} \left(f_{\left[\begin{smallmatrix} 0 \\ 1/6 \end{smallmatrix} \right]}(\tau) + f_{\left[\begin{smallmatrix} 2/6 \\ 1/6 \end{smallmatrix} \right]}(\tau) + f_{\left[\begin{smallmatrix} 2/6 \\ 3/6 \end{smallmatrix} \right]}(\tau) + f_{\left[\begin{smallmatrix} 2/6 \\ 5/6 \end{smallmatrix} \right]}(\tau) \right) \\
&= \frac{1}{8} \sum_{(a,b) \in T} f_{\left[\begin{smallmatrix} a/6 \\ b/6 \end{smallmatrix} \right]}(\tau) - \frac{1}{8} f_{\left[\begin{smallmatrix} 0 \\ 3/6 \end{smallmatrix} \right]}(\tau) \\
&= \frac{g_2(\tau)g_3(\tau)}{8\Delta(\tau)} \sum_{(a,b) \in T} \wp \left(\frac{a}{6}\tau + \frac{b}{6}; [\tau, 1] \right) - \frac{1}{8} f_{\left[\begin{smallmatrix} 0 \\ 3/6 \end{smallmatrix} \right]}(\tau) \\
&= \frac{g_2(\tau)g_3(\tau)}{8\Delta(\tau)} \sum_{(a,b) \in T} \left(\sum_{(m,n) \in \mathbb{Z}^2} \frac{36}{\{(a-6m)\tau + (b-6n)\}^2} - \sum_{(m,n) \in \mathbb{Z}^2 \setminus \{(0,0)\}} \frac{1}{(m\tau + n)^2} \right) \\
&\quad - \frac{1}{8} f_{\left[\begin{smallmatrix} 0 \\ 3/6 \end{smallmatrix} \right]}(\tau) \\
&= \frac{9g_2(\tau)g_3(\tau)}{8\Delta(\tau)} \left(\sum_{(m,n) \in \mathbb{Z}^2} \frac{4}{\{-2m\tau + (1-2n)\}^2} - \sum_{(m,n) \in \mathbb{Z}^2 \setminus \{(0,0)\}} \frac{1}{(m\tau + n)^2} \right) \\
&\quad - \frac{1}{8} f_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix} \right]}(\tau) \\
&= \frac{9g_2(\tau)g_3(\tau)}{8\Delta(\tau)} \wp \left(\frac{1}{2}; [\tau, 1] \right) - \frac{1}{8} f_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix} \right]}(\tau)
\end{aligned}$$

$$= f_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix} \right]}(\tau). \quad \square$$

Lemma 3.10. *We have*

$$f_{\left[\begin{smallmatrix} 0 \\ 1/3 \end{smallmatrix} \right]}(\tau) = \frac{1}{3} \left(f_{\left[\begin{smallmatrix} 0 \\ 1/6 \end{smallmatrix} \right]}(\tau) + f_{\left[\begin{smallmatrix} 3/6 \\ 1/6 \end{smallmatrix} \right]}(\tau) + f_{\left[\begin{smallmatrix} 3/6 \\ 4/6 \end{smallmatrix} \right]}(\tau) \right).$$

Proof. Let $T = \{(0, 1), (0, 4), (3, 1), (3, 4)\}$. We achieve by Proposition 2.3(i) and the definitions (2) and (4) that

$$\begin{aligned} & \frac{1}{3} \left(f_{\left[\begin{smallmatrix} 0 \\ 1/6 \end{smallmatrix} \right]}(\tau) + f_{\left[\begin{smallmatrix} 3/6 \\ 1/6 \end{smallmatrix} \right]}(\tau) + f_{\left[\begin{smallmatrix} 3/6 \\ 4/6 \end{smallmatrix} \right]}(\tau) \right) \\ &= \frac{1}{3} \sum_{(a,b) \in T} f_{\left[\begin{smallmatrix} a/6 \\ b/6 \end{smallmatrix} \right]}(\tau) - \frac{1}{3} f_{\left[\begin{smallmatrix} 0 \\ 4/6 \end{smallmatrix} \right]}(\tau) \\ &= \frac{g_2(\tau)g_3(\tau)}{3\Delta(\tau)} \sum_{(a,b) \in T} \wp \left(\frac{a}{6}\tau + \frac{b}{6}; [\tau, 1] \right) - \frac{1}{3} f_{\left[\begin{smallmatrix} 0 \\ 1/3 \end{smallmatrix} \right]}(\tau) \\ &= \frac{g_2(\tau)g_3(\tau)}{3\Delta(\tau)} \sum_{(a,b) \in T} \left(\sum_{(m,n) \in \mathbb{Z}^2} \frac{36}{\{(a-6m)\tau + (b-6n)\}^2} - \sum_{(m,n) \in \mathbb{Z}^2 \setminus \{(0,0)\}} \frac{1}{(m\tau + n)^2} \right) \\ & \quad - \frac{1}{3} f_{\left[\begin{smallmatrix} 0 \\ 1/3 \end{smallmatrix} \right]}(\tau) \\ &= \frac{4g_2(\tau)g_3(\tau)}{3\Delta(\tau)} \left(\sum_{(m,n) \in \mathbb{Z}^2} \frac{9}{\{-3m\tau + (1-3n)\}^2} - \sum_{(m,n) \in \mathbb{Z}^2 \setminus \{(0,0)\}} \frac{1}{(m\tau + n)^2} \right) \\ & \quad - \frac{1}{3} f_{\left[\begin{smallmatrix} 0 \\ 1/3 \end{smallmatrix} \right]}(\tau) \\ &= \frac{4g_2(\tau)g_3(\tau)}{3\Delta(\tau)} \wp \left(\frac{1}{3}; [\tau, 1] \right) - \frac{1}{3} f_{\left[\begin{smallmatrix} 0 \\ 1/3 \end{smallmatrix} \right]}(\tau) \\ &= f_{\left[\begin{smallmatrix} 0 \\ 1/3 \end{smallmatrix} \right]}(\tau). \quad \square \end{aligned}$$

Lemma 3.11. *We have the following identities:*

- (i) $g_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix} \right]}(\tau)^{12} g_{\left[\begin{smallmatrix} 1/2 \\ 0 \end{smallmatrix} \right]}(\tau)^{12} g_{\left[\begin{smallmatrix} 1/2 \\ 1/2 \end{smallmatrix} \right]}(\tau)^{12} = -2^{12}$.
- (ii) $g_{\left[\begin{smallmatrix} 0 \\ 1/3 \end{smallmatrix} \right]}(\tau)^{12} g_{\left[\begin{smallmatrix} 1/3 \\ 0 \end{smallmatrix} \right]}(\tau)^{12} g_{\left[\begin{smallmatrix} 1/3 \\ 1/3 \end{smallmatrix} \right]}(\tau)^{12} g_{\left[\begin{smallmatrix} 1/3 \\ 2/3 \end{smallmatrix} \right]}(\tau)^{12} = 3^6$.

Proof. For convenience, let

$$G_2(\tau) = g_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix} \right]}(\tau)^{12} g_{\left[\begin{smallmatrix} 1/2 \\ 0 \end{smallmatrix} \right]}(\tau)^{12} g_{\left[\begin{smallmatrix} 1/2 \\ 1/2 \end{smallmatrix} \right]}(\tau)^{12},$$

$$G_3(\tau) = g_{\left[\begin{smallmatrix} 0 \\ 1/3 \end{smallmatrix} \right]}(\tau)^{12} g_{\left[\begin{smallmatrix} 1/3 \\ 0 \end{smallmatrix} \right]}(\tau)^{12} g_{\left[\begin{smallmatrix} 1/3 \\ 1/3 \end{smallmatrix} \right]}(\tau)^{12} g_{\left[\begin{smallmatrix} 1/3 \\ 2/3 \end{smallmatrix} \right]}(\tau)^{12}.$$

Let $N \in \{2, 3\}$. By the definition (5), the leading coefficient of $G_N(\tau)$ is given by

$$\begin{cases} -2^{12} & \text{if } N = 2, \\ 3^6 & \text{if } N = 3. \end{cases}$$

One can readily show by Proposition 2.3(i)–(iii) that $G_N(\tau)$ is invariant under the action of $\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})/\{\pm I_2\}$, which implies that $G_N(\tau)$ belongs to \mathcal{F}_1 . Since the width of $i\infty$ on the modular curve $X(1)$ is 1, we see by Proposition 2.4(i) that

$$\begin{aligned} \mathrm{ord}_{i\infty} G_N(\tau) &= \mathrm{ord}_q G_N(\tau) \\ &= \begin{cases} \left\{ 12 \cdot \frac{1}{2} (\mathbf{B}_2(0) + 2\mathbf{B}_2(1/2)) \right\} & \text{if } N = 2, \\ \left\{ 12 \cdot \frac{1}{2} (\mathbf{B}_2(0) + 3\mathbf{B}_2(1/3)) \right\} & \text{if } N = 3 \end{cases} \\ &= 0. \end{aligned}$$

Thus $G_N(\tau)$ is holomorphic functions on $X(1)$; and hence it is a constant, namely,

$$G_2(\tau) = -2^{12} \quad \text{and} \quad G_3(\tau) = 3^6. \quad \square$$

4. Class fields over imaginary quadratic fields

In this section, we shall briefly review an explicit version of Shimura's reciprocity law.

Let K be an imaginary quadratic field of discriminant d_K , and let \mathcal{O}_K be its ring of integers. For a nontrivial ideal \mathfrak{m} of \mathcal{O}_K , let $K_{\mathfrak{m}}$ denote the ray class field of K modulo \mathfrak{m} . In particular, $K_{\mathcal{O}_K}$ becomes the maximal unramified abelian extension of K , called the Hilbert class field of K and denoted by H_K . For more details, we refer to [2, §5 and §8] or [7, Chapter V].

Lemma 4.1. *We have the degree formula*

$$[K_{\mathfrak{m}} : H_K] = \frac{\phi(\mathfrak{m})\omega(\mathfrak{m})}{\omega_K},$$

where ϕ is the multiplicative Euler function for ideals given by

$$\phi(\mathfrak{p}^n) = (N_{K/\mathbb{Q}}(\mathfrak{p}) - 1)N_{K/\mathbb{Q}}(\mathfrak{p})^{n-1}$$

for each prime ideal power \mathfrak{p}^n ($n \geq 1$), $\omega(\mathfrak{m})$ is the number of units in \mathcal{O}_K which are congruent to 1 modulo \mathfrak{m} , and $\omega_K = \omega(\mathcal{O}_K)$.

Proof. See [16, Theorem 1 in Chapter VI]. □

Throughout this section, we let N be a positive integer and $\mathfrak{m} = N\mathcal{O}_K$. Further, we fix an element τ_K of \mathbb{H} as

$$(7) \quad \tau_K = \begin{cases} \sqrt{d_K}/2 & \text{if } d_K \equiv 0 \pmod{4}, \\ (-1 + \sqrt{d_K})/2 & \text{if } d_K \equiv 1 \pmod{4}; \end{cases}$$

and hence $\mathcal{O}_K = \mathbb{Z}[\tau_K]$ ([2, p. 103]).

Proposition 4.2. *Let $E = E_{\mathcal{O}_K}$ be the elliptic curve with parametrization $\varphi_E : \mathbb{C}/\mathcal{O}_K \rightarrow E$ given in §2, and let h_E be the Weber function on E defined in (1).*

- (i) $H_K = K(j(E)) = K(j(\tau_K))$.
- (ii) $K_{\mathfrak{m}} = K(f(\tau_K) \mid f \in \mathcal{F}_N \text{ is finite at } \tau_K)$.
- (iii) *If $N \geq 2$, then $K_{\mathfrak{m}} = H_K(h_E(\varphi_E(1/N)))$.*
- (iv) *If $K \neq \mathbb{Q}(\sqrt{-1}), \mathbb{Q}(\sqrt{-3})$ and $N \geq 2$, then*

$$K_{\mathfrak{m}} = K \left(g_{\begin{bmatrix} 0 \\ 1/N \end{bmatrix}}(\tau_K)^{12Nn/\gcd(6, N)} \right)$$

for any nonzero integer n .

Proof. (i) See [15, Theorem 1 in Chapter 10].

(ii) See [15, Corollary to Theorem 2 in Chapter 10].

(iii) See [15, Corollary to Theorem 7 in Chapter 10].

(iv) See [8, Theorem 4.2]. □

Remark 4.3. If $K \neq \mathbb{Q}(\sqrt{-1}), \mathbb{Q}(\sqrt{-3})$ and $N \geq 2$, then we get by the definitions (1) and (4) that

$$h_E(\varphi_E(1/N)) = f_{\begin{bmatrix} 0 \\ 1/N \end{bmatrix}}(\tau_K).$$

Let $\min(\tau_K, \mathbb{Q}) = x^2 + bx + c \in \mathbb{Z}[x]$, and let

$$W_{K, N} = \left\{ \begin{bmatrix} t - bs & -cs \\ s & t \end{bmatrix} \in \mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z}) \mid t, s \in (\mathbb{Z}/N\mathbb{Z}) \right\}$$

which is a subgroup of $\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$.

Proposition 4.4 (Shimura's reciprocity law). *There is a surjection*

$$\begin{aligned} W_{K, N} &\rightarrow \mathrm{Gal}(K_{\mathfrak{m}}/H_K) \\ \gamma &\mapsto (f(\tau_K) \mapsto f^\gamma(\tau_K) \mid f \in \mathcal{F}_N \text{ is finite at } \tau_K) \end{aligned}$$

whose kernel is

$$\begin{cases} \left\{ \pm \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \pm \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \right\} & \text{if } K = \mathbb{Q}(\sqrt{-1}), \\ \left\{ \pm \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \pm \begin{bmatrix} 1 & 1 \\ -1 & 0 \end{bmatrix}, \pm \begin{bmatrix} 0 & -1 \\ 1 & 1 \end{bmatrix} \right\} & \text{if } K = \mathbb{Q}(\sqrt{-3}), \\ \left\{ \pm \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right\} & \text{if } K \neq \mathbb{Q}(\sqrt{-1}), \mathbb{Q}(\sqrt{-3}). \end{cases}$$

Proof. See [22, §3]. □

Let $C(d_K)$ be the form class group of properly equivalence classes of primitive positive definite quadratic forms of discriminant d_K under Dirichlet composition. See [2, Theorem 3.9] for the details. Note that every element of $C(d_K)$

is uniquely represented by a reduced form $Q = ax^2 + bxy + cy^2 \in \mathbb{Z}[x, y]$ characterized by the condition

$$(8) \quad (-a < b \leq a < c \text{ or } 0 \leq b \leq a = c) \quad \text{and} \quad b^2 - 4ac = d_K$$

([2, Theorem 2.8]). For each reduced form $Q = ax^2 + bxy + cy^2$, let

$$(9) \quad \tau_Q = \frac{-b + \sqrt{d_K}}{2a} \in \mathbb{H}.$$

If $d_K \equiv 0 \pmod{4}$, then for each prime p we set

$$(10) \quad \gamma_{Q,p} = \begin{cases} \begin{bmatrix} a & b/2 \\ 0 & 1 \end{bmatrix} & \text{if } p \nmid a, \\ \begin{bmatrix} -b/2 & -c \\ 1 & 0 \end{bmatrix} & \text{if } p \mid a \text{ and } p \nmid c, \\ \begin{bmatrix} -a - b/2 & -c - b/2 \\ 1 & -1 \end{bmatrix} & \text{if } p \mid a \text{ and } p \mid c. \end{cases}$$

Otherwise, if $d_K \equiv 1 \pmod{4}$, then we set

$$(11) \quad \gamma_{Q,p} = \begin{cases} \begin{bmatrix} a & (b-1)/2 \\ 0 & 1 \end{bmatrix} & \text{if } p \nmid a, \\ \begin{bmatrix} -(b+1)/2 & -c \\ 1 & 0 \end{bmatrix} & \text{if } p \mid a \text{ and } p \nmid c, \\ \begin{bmatrix} -a - (b+1)/2 & -c + (1-b)/2 \\ 1 & -1 \end{bmatrix} & \text{if } p \mid a \text{ and } p \mid c. \end{cases}$$

Then, there exists a unique matrix r_Q in $\text{GL}_2(\mathbb{Z}/N\mathbb{Z})$ satisfying $r_Q \equiv r_{Q,p} \pmod{N \cdot \mathbb{Z}_p}$ for all primes p by the Chinese remainder theorem.

Proposition 4.5 (Shimura's reciprocity law). *We have an injective map*

$$\begin{aligned} \mathcal{C}(d_K) &\rightarrow \text{Gal}(K_m/K) \\ \mathcal{C}^{-1} &\mapsto (f(\tau_K) \mapsto f^{\gamma_Q}(\tau_Q) \mid f \in \mathcal{F}_N \text{ is finite at } \tau_K). \end{aligned}$$

Here, Q^{-1} is a reduced form which is the inverse of Q . And, the restriction to H_K followed by the above map gives rise to an isomorphism of $\mathcal{C}(d_K)$ onto $\text{Gal}(H_K/K)$.

Proof. See [22, §6]. □

Remark 4.6. (i) The identity of $\mathcal{C}(d_K)$ is represented by the reduced form

$$\begin{cases} x^2 - \frac{d_K}{4}y^2 & \text{if } d_K \equiv 0 \pmod{4}, \\ x^2 + xy + \frac{1-d_K}{4}y^2 & \text{if } d_K \equiv 1 \pmod{4} \end{cases}$$

([2, Theorem 3.9]). One can readily show by the definitions (7), (9), (10) and (11) that the identity of $\mathcal{C}(d_K)$ corresponds to the identity of $\text{Gal}(K_m/K)$ via the map given in Proposition 4.5.

(ii) Let $Q = ax^2 + bxy + cy^2$ be a reduced form of discriminant d_K . Then, the condition (8) implies

$$a \leq \sqrt{\frac{|d_K|}{3}}$$

([2, p. 29]). Moreover, if Q is a nonidentity, then we get $a \geq 2$.

(iii) Let h_K be the class number of K . It is well known that

$$h_K = 1 \iff d_K = -3, -4, -7, -8, -11, -19, -43, -67, -163$$

due to Heegner ([6]), Baker ([1]) and Stark ([20] and [21]).

(iv) Let $N \geq 2$. If $K = \mathbb{Q}(\sqrt{-1})$ or $\mathbb{Q}(\sqrt{-3})$, then we attain by (iii) and Proposition 4.2(iii) that $H_K = K$ and

$$K_{\mathfrak{m}} = K(h_E(\varphi_E(1/N))).$$

5. Generation of ray class fields modulo 2

We are ready to prove our main theorems. Henceforth we assume that K is an imaginary quadratic field different from $\mathbb{Q}(\sqrt{-1})$ and $\mathbb{Q}(\sqrt{-3})$. Furthermore, we let $E = E_{\mathcal{O}_K}$ be the elliptic curve with parametrization $\varphi_E : \mathbb{C}/\mathcal{O}_K \rightarrow E$ stated in §2, and let h_E be the Weber function on E .

Let $\mathfrak{m} = 2\mathcal{O}_K$. By Proposition 4.4 we have

$$(12) \quad \begin{aligned} \text{Gal}(K_{\mathfrak{m}}/H_K) &\simeq W_{K,2}/\{\pm I_2\} \\ &= \begin{cases} \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \right\} & \text{if } d_K \equiv 0 \pmod{8}, \\ \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \right\} & \text{if } d_K \equiv 4 \pmod{8}, \\ \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix} \right\} & \text{if } d_K \equiv 5 \pmod{8}, \\ \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right\} & \text{if } d_K \equiv 1 \pmod{8}. \end{cases} \end{aligned}$$

Let $E = E_{\mathcal{O}_K}$ be the elliptic curve with parametrization $\varphi_E : \mathbb{C}/\mathcal{O}_K \rightarrow E$ stated in §2, and let h_E be the Weber function on E .

Theorem 5.1. *Suppose that 2 does not split in K . Then, $h_E(\varphi_E(1/2))$ generates $K_{\mathfrak{m}}$ over K .*

Proof. Since 2 is ramified or inert in K , we get $d_K \equiv 0 \pmod{4}$ or $d_K \equiv 5 \pmod{8}$. And, it follows from (12) and Proposition 2.3(iii) that the Galois

conjugates of $f_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix}\right]}(\tau_K)$ over H_K are given by

$$\begin{cases} f_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix}\right]}(\tau_K), f_{\left[\begin{smallmatrix} 1/2 \\ 1/2 \end{smallmatrix}\right]}(\tau_K) & \text{if } d_K \equiv 0 \pmod{8}, \\ f_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix}\right]}(\tau_K), f_{\left[\begin{smallmatrix} 1/2 \\ 0 \end{smallmatrix}\right]}(\tau_K) & \text{if } d_K \equiv 4 \pmod{8}, \\ f_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix}\right]}(\tau_K), f_{\left[\begin{smallmatrix} 1/2 \\ 0 \end{smallmatrix}\right]}(\tau_K), f_{\left[\begin{smallmatrix} 1/2 \\ 1/2 \end{smallmatrix}\right]}(\tau_K) & \text{if } d_K \equiv 5 \pmod{8}. \end{cases}$$

Let

$$F = K(h_E(\varphi_E(1/2))) = K\left(f_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix}\right]}(\tau_K)\right)$$

which is a subfield of K_m by Proposition 4.2(ii). Now that F is an abelian extension of K , it contains all Galois conjugates of $f_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix}\right]}(\tau_K)$ over K . Furthermore, since

$$f_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix}\right]}(\tau_K) + f_{\left[\begin{smallmatrix} 1/2 \\ 0 \end{smallmatrix}\right]}(\tau_K) + f_{\left[\begin{smallmatrix} 1/2 \\ 1/2 \end{smallmatrix}\right]}(\tau_K) = 0$$

by Lemma 3.2, F contains all three values $f_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix}\right]}(\tau_K)$, $f_{\left[\begin{smallmatrix} 1/2 \\ 0 \end{smallmatrix}\right]}(\tau_K)$ and $f_{\left[\begin{smallmatrix} 1/2 \\ 1/2 \end{smallmatrix}\right]}(\tau_K)$. Thus we achieve by Lemma 3.2 that

$$\begin{aligned} F &\ni \frac{f_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix}\right]}(\tau_K)f_{\left[\begin{smallmatrix} 1/2 \\ 0 \end{smallmatrix}\right]}(\tau_K)f_{\left[\begin{smallmatrix} 1/2 \\ 1/2 \end{smallmatrix}\right]}(\tau_K)}{f_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix}\right]}(\tau_K)f_{\left[\begin{smallmatrix} 1/2 \\ 0 \end{smallmatrix}\right]}(\tau_K) + f_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix}\right]}(\tau_K)f_{\left[\begin{smallmatrix} 1/2 \\ 1/2 \end{smallmatrix}\right]}(\tau_K) + f_{\left[\begin{smallmatrix} 1/2 \\ 0 \end{smallmatrix}\right]}(\tau_K)f_{\left[\begin{smallmatrix} 1/2 \\ 1/2 \end{smallmatrix}\right]}(\tau_K)} \\ &= \frac{j(\tau_K)(j(\tau_K) - 1728)^2}{-27 \cdot 1728 j(\tau_K)(j(\tau_K) - 1728)} \\ &= \frac{-j(\tau_K) + 1728}{27 \cdot 1728}. \end{aligned}$$

Therefore, we conclude by Proposition 4.2(i) and (iii) that $F = K_m$. This proves the theorem. \square

Lemma 5.2. *Let $u + vi \in \mathbb{H}$, and let $t = e^{-2\pi v}$.*

- (i) $\left|g_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix}\right]}(u + vi)\right| \leq 2t^{1/12}e^{-2+2/(1-t)}.$
- (ii) $\left|g_{\left[\begin{smallmatrix} 1/2 \\ s \end{smallmatrix}\right]}(u + vi)\right| \leq t^{-1/24}e^{t^{1/2}(1+2/(1-t))}$ for any $s \in \mathbb{Q}$.

Proof. (i) We derive that

$$\begin{aligned} \left|g_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix}\right]}(u + vi)\right| &\leq 2t^{(1/2)\mathbf{B}_2(0)} \prod_{n=1}^{\infty} (1 + t^n)^2 \quad \text{by the definition (5)} \\ &\leq 2t^{1/12} \prod_{n=1}^{\infty} e^{2t^n} \quad \text{because } 1 + x < e^x \text{ for all } x > 0 \end{aligned}$$

$$= 2t^{1/12}e^{-2+2/(1-t)}.$$

(ii) We see that

$$\begin{aligned} & \left| g_{\left[\begin{smallmatrix} 1/2 \\ s \end{smallmatrix} \right]}(u + vi) \right| \\ & \leq t^{(1/2)\mathbf{B}_2(1/2)}(1+t^{1/2}) \prod_{n=1}^{\infty} (1+t^{n+1/2})(1+t^{n-1/2}) \quad \text{by the definition (5)} \\ & \leq t^{-1/24}(1+t^{1/2}) \prod_{n=1}^{\infty} (1+t^{n-1/2})^2 \\ & \leq t^{-1/24}e^{t^{1/2}} \prod_{n=1}^{\infty} e^{2t^{n-1/2}} \quad \text{since } 1+x < e^x \text{ for all } x > 0 \\ & = t^{-1/24}e^{t^{1/2}(1+2/(1-t))}. \quad \square \end{aligned}$$

Theorem 5.3. *If 2 splits in K , then $h_E(\varphi_E(1/2))$ generates K_m over K .*

Proof. Since 2 splits in K , we have $d_K \equiv 1 \pmod{8}$ and $K_m = H_K$ by Lemma 4.1 or (12). If $h_K = 1$, then the theorem is obvious. So, we let $h_K \geq 2$, and hence $d_K \leq -15$ by Remark 4.6(iii).

Suppose on the contrary that $h_E(\varphi_E(1/2)) = f_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix} \right]}(\tau_K)$ does not generate H_K over K . Then, by Proposition 4.5, there is a nonidentity reduced form $Q = ax^2 + bxy + cy^2$ of discriminant d_K which leaves $f_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix} \right]}(\tau_K)$ fixed. Let

$$f = f_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix} \right]}(\tau_K), \quad g = g_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix} \right]}(\tau_K)^{12} \quad \text{and} \quad h = g^{Q^{-1}}.$$

Then we get by Propositions 4.5 and 2.3(iii) that

$$h = \left(g_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix} \right]}(\tau)^{12} \right)^{\gamma_Q} (\tau_Q) = g_{\left[\begin{smallmatrix} r \\ s \end{smallmatrix} \right]} \left(\frac{-b + \sqrt{d_K}}{2a} \right)^{12} \quad \text{for some } \begin{bmatrix} r \\ s \end{bmatrix} \in (1/2)\mathbb{Z}^2 \setminus \mathbb{Z}^2.$$

Now, we establish that

$$\begin{aligned} |g| & \leq 2^{12}e^{-\pi\sqrt{|d_K|}}e^{-24+24/(1-e^{-\pi\sqrt{|d_K|}})} \quad \text{by Lemma 5.2(i)} \\ & \leq 2^{12}e^{-\pi\sqrt{|d_K|}}e^{-24+24/(1-e^{-\pi\sqrt{15}})} \quad \text{by the fact } d_K \leq -15 \\ & < 4097e^{-\pi\sqrt{|d_K|}} \end{aligned}$$

and

$$\begin{aligned} |h| & \leq \begin{cases} 2^{12}e^{-\pi\sqrt{|d_K|}/a}e^{-24+24/(1-e^{-\pi\sqrt{|d_K|}/a})} & \text{if } \langle r \rangle = 0, \\ e^{\pi\sqrt{|d_K|}/2a}e^{12e^{-\pi\sqrt{|d_K|}/2a}(1+2/(1-e^{-\pi\sqrt{|d_K|}/a}))} & \text{if } \langle r \rangle = 1/2 \end{cases} \\ & \quad \text{by Lemma 5.2(ii)} \\ & \leq \begin{cases} 2^{12}e^{-\pi\sqrt{3}}e^{-24+24/(1-e^{-\pi\sqrt{3}})} & \text{if } \langle r \rangle = 0, \\ e^{\pi\sqrt{|d_K|}/4}e^{12e^{-\pi\sqrt{3}/2}(1+2/(1-e^{-\pi\sqrt{3}}))} & \text{if } \langle r \rangle = 1/2, \end{cases} \end{aligned}$$

$$\begin{aligned}
& \text{because } \sqrt{3} \leq \sqrt{|d_K|}/a \text{ by Remark 4.6(ii)} \\
& < \begin{cases} 20 & \text{if } \langle r \rangle = 0, \\ 11e^{\pi\sqrt{|d_K|}/4} & \text{if } \langle r \rangle = 1/2 \end{cases} \\
& \leq 11e^{\pi\sqrt{|d_K|}/4}.
\end{aligned}$$

We obtain by the fact $f = f^{Q^{-1}}$ and Lemma 3.6 that

$$\frac{g^3 + 72g^2 + 384g - 8192}{g} = \frac{h^3 + 72h^2 + 384h - 8192}{h},$$

from which we attain

$$(g - h)(g^2h + gh^2 + 72gh + 8192) = 0.$$

Since g generates K_m over K by Proposition 4.2(iv), we see that $g \neq h$; and hence

$$g^2h + gh^2 + 72gh + 8192 = 0.$$

We then derive that

$$\begin{aligned}
8192 &= |g^2h + gh^2 + 72gh| \\
&\leq |g^2h| + |gh^2| + 72|gh| \\
&< \left(4097e^{-\pi\sqrt{|d_K|}}\right)^2 \left(11e^{\pi\sqrt{|d_K|}/4}\right) + \left(4097e^{-\pi\sqrt{|d_K|}}\right) \left(11e^{\pi\sqrt{|d_K|}/4}\right)^2 \\
&\quad + 72 \left(4097e^{-\pi\sqrt{|d_K|}}\right) \left(11e^{\pi\sqrt{|d_K|}/4}\right) \\
&= 4097^2 \cdot 11e^{-7\pi\sqrt{|d_K|}/4} + 4097 \cdot 11^2 e^{-\pi\sqrt{|d_K|}/2} \\
&\quad + 72 \cdot 4097 \cdot 11e^{-3\pi\sqrt{|d_K|}/4} \\
&< 1484 \quad \text{because } |d_K| \geq 15,
\end{aligned}$$

which yields a contradiction. As for the last inequality in the above, we rely on evaluation using Maple 2016 software. Therefore, we conclude that $h_E(\varphi_E(1/2))$ generates $K_m = H_K$ over K . \square

6. Generation of ray class fields modulo 3

Let $\mathfrak{m} = 3\mathcal{O}_K$. By Proposition 4.4 we know that

$$\text{Gal}(K_m/H_K) \simeq W_{K,3}/\{\pm I_2\}$$

$$(13) \quad = \begin{cases} \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}, \begin{bmatrix} 2 & 0 \\ 1 & 2 \end{bmatrix} \right\} & \text{if } d_K \equiv 0 \pmod{12}, \\ \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 2 & 2 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 2 \\ 1 & 1 \end{bmatrix} \right\} & \text{if } d_K \equiv 9 \pmod{12}, \\ \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 1 & 1 \end{bmatrix}, \begin{bmatrix} 2 & 2 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 0 & 2 \\ 1 & 0 \end{bmatrix} \right\} & \text{if } d_K \equiv 8 \pmod{12}, \\ \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 2 & 1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix} \right\} & \text{if } d_K \equiv 5 \pmod{12}, \\ \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \right\} & \text{if } d_K \equiv 4 \pmod{12}, \\ \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 1 & 2 \end{bmatrix} \right\} & \text{if } d_K \equiv 1 \pmod{12}. \end{cases}$$

Theorem 6.1. *Suppose that 3 does not split in K . Then, $h_E(\varphi_E(1/3))$ generates K_m over K .*

Proof. Since 3 is ramified or inert in K , we have $d_K \equiv 0$ or $2 \pmod{3}$. It follows from (13) and Proposition 2.3(iii) that the Galois conjugates of $f_{\begin{bmatrix} 0 \\ 1/3 \end{bmatrix}}(\tau_K)$ over H_K are given by

$$\begin{cases} f_{\begin{bmatrix} 0 \\ 1/3 \end{bmatrix}}(\tau_K), f_{\begin{bmatrix} 1/3 \\ 1/3 \end{bmatrix}}(\tau_K), f_{\begin{bmatrix} 1/3 \\ 2/3 \end{bmatrix}}(\tau_K) & \text{if } d_K \equiv 0 \pmod{12}, \\ f_{\begin{bmatrix} 0 \\ 1/3 \end{bmatrix}}(\tau_K), f_{\begin{bmatrix} 1/3 \\ 0 \end{bmatrix}}(\tau_K), f_{\begin{bmatrix} 1/3 \\ 1/3 \end{bmatrix}}(\tau_K) & \text{if } d_K \equiv 9 \pmod{12}, \\ f_{\begin{bmatrix} 0 \\ 1/3 \end{bmatrix}}(\tau_K), f_{\begin{bmatrix} 1/3 \\ 0 \end{bmatrix}}(\tau_K), f_{\begin{bmatrix} 1/3 \\ 1/3 \end{bmatrix}}(\tau_K), f_{\begin{bmatrix} 1/3 \\ 2/3 \end{bmatrix}}(\tau_K) & \text{if } d_K \equiv 8 \text{ or } 5 \pmod{12}. \end{cases}$$

Let

$$F = K(h_E(\varphi_E(1/3))) = K\left(f_{\begin{bmatrix} 0 \\ 1/3 \end{bmatrix}}(\tau_K)\right)$$

which is an abelian extension of K . Since

$$f_{\begin{bmatrix} 0 \\ 1/3 \end{bmatrix}}(\tau) + f_{\begin{bmatrix} 1/3 \\ 0 \end{bmatrix}}(\tau) + f_{\begin{bmatrix} 1/3 \\ 1/3 \end{bmatrix}}(\tau) + f_{\begin{bmatrix} 1/3 \\ 2/3 \end{bmatrix}}(\tau) = 0$$

by Lemma 3.7, F contains all four distinct values $f_{\begin{bmatrix} 0 \\ 1/3 \end{bmatrix}}(\tau_K)$, $f_{\begin{bmatrix} 1/3 \\ 0 \end{bmatrix}}(\tau_K)$, $f_{\begin{bmatrix} 1/3 \\ 1/3 \end{bmatrix}}(\tau_K)$ and $f_{\begin{bmatrix} 1/3 \\ 2/3 \end{bmatrix}}(\tau_K)$. We then deduce that

$$F \ni \frac{\left(f_{\begin{bmatrix} 1/3 \\ 1/3 \end{bmatrix}}(\tau_K) - f_{\begin{bmatrix} 1/3 \\ 0 \end{bmatrix}}(\tau_K)\right)^{24} \left(f_{\begin{bmatrix} 1/3 \\ 2/3 \end{bmatrix}}(\tau_K) - f_{\begin{bmatrix} 1/3 \\ 1/3 \end{bmatrix}}(\tau_K)\right)^{12}}{\left(f_{\begin{bmatrix} 1/3 \\ 1/3 \end{bmatrix}}(\tau_K) - f_{\begin{bmatrix} 0 \\ 1/3 \end{bmatrix}}(\tau_K)\right)^{24} \left(f_{\begin{bmatrix} 1/3 \\ 1/3 \end{bmatrix}}(\tau_K) - f_{\begin{bmatrix} 0 \\ 1/3 \end{bmatrix}}(\tau_K)\right)^{12}}$$

$$\begin{aligned}
&= \frac{g\left[\begin{smallmatrix} 0 \\ 1/3 \end{smallmatrix}\right](\tau_K)^{108}}{g\left[\begin{smallmatrix} 1/3 \\ 0 \end{smallmatrix}\right](\tau_K)^{36}g\left[\begin{smallmatrix} 1/3 \\ 1/3 \end{smallmatrix}\right](\tau_K)^{36}g\left[\begin{smallmatrix} 1/3 \\ 2/3 \end{smallmatrix}\right](\tau_K)^{36}} \\
&\quad \text{by Propositions 2.3(i) and 2.4(ii)} \\
&= 3^{-18}g\left[\begin{smallmatrix} 0 \\ 1/3 \end{smallmatrix}\right](\tau_K)^{144} \quad \text{by Lemma 3.11.}
\end{aligned}$$

Therefore, we conclude by Proposition 4.2(iv) that $F = K_m$ as desired. \square

Lemma 6.2. *Let $u + vi \in \mathbb{H}$ with $t = e^{-2\pi v}$, and let $\begin{bmatrix} r \\ s \end{bmatrix} \in (1/3)\mathbb{Z}^2 \setminus \mathbb{Z}^2$. If $t \leq e^{-\pi\sqrt{3}}$, then we have*

$$\begin{aligned}
\left|g\left[\begin{smallmatrix} r \\ s \end{smallmatrix}\right](u + vi)\right| &< \begin{cases} 1.75t^{1/12} & \text{if } \langle r \rangle = 0, \\ 1.25t^{-1/36} & \text{if } \langle r \rangle \neq 0 \end{cases} \\
&\leq 1.25t^{-1/36}.
\end{aligned}$$

Proof. We may assume that $\begin{bmatrix} r \\ s \end{bmatrix} \in \left\{\begin{bmatrix} 0 \\ 1/3 \end{bmatrix}, \begin{bmatrix} 1/3 \\ 0 \end{bmatrix}, \begin{bmatrix} 1/3 \\ 1/3 \end{bmatrix}, \begin{bmatrix} 1/3 \\ 2/3 \end{bmatrix}\right\}$ by Proposition 2.3(i). Then we derive that

$$\begin{aligned}
\left|g\left[\begin{smallmatrix} r \\ s \end{smallmatrix}\right](u + vi)\right| &\leq \begin{cases} \sqrt{3}t^{(1/2)\mathbf{B}_2(0)} \prod_{n=1}^{\infty} (1 + t^n)^2 & \text{if } r = 0, \\ t^{(1/2)\mathbf{B}_2(1/3)} (1 + t^{1/3}) \prod_{n=1}^{\infty} (1 + t^{n-1/3})^2 & \text{if } r = 1/3, \end{cases} \\
&\quad \text{by the definition (5)} \\
&\leq \begin{cases} \sqrt{3}t^{1/12} \prod_{n=1}^{\infty} e^{2t^n} & \text{if } r = 0, \\ t^{-1/36} e^{t^{1/3}} \prod_{n=1}^{\infty} e^{2t^{n-1/3}} & \text{if } r = 1/3, \end{cases} \\
&\quad \text{since } 1 + x < e^x \text{ for all } x > 0 \\
&= \begin{cases} \sqrt{3}t^{1/12} e^{2t/(1-t)} & \text{if } r = 0, \\ t^{-1/36} e^{t^{1/3} + 2t^{2/3}/(1-t)} & \text{if } r = 1/3. \end{cases}
\end{aligned}$$

Moreover, now that $t \leq e^{-\pi\sqrt{3}}$, we get that

$$\begin{aligned}
\left|g\left[\begin{smallmatrix} r \\ s \end{smallmatrix}\right](u + vi)\right| &< \begin{cases} 1.75t^{1/12} & \text{if } r = 0, \\ 1.25t^{-1/36} & \text{if } r = 1/3 \end{cases} \\
&\leq 1.25t^{-1/36}. \quad \square
\end{aligned}$$

Lemma 6.3. *Suppose that $h_K \geq 2$ and $d_K \leq -20$. Let $Q = ax^2 + bxy + cy^2$ be a nonidentity reduced form of discriminant d_K , and let $\gamma \in \text{GL}_2(\mathbb{Z}/3\mathbb{Z})$.*

$$(i) \quad \left|g\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix}\right](\tau_Q)\right| > 1.98e^{-\pi\sqrt{|d_K|}/24}.$$

$$(ii) \left| \frac{j(\tau_Q)^2(j(\tau_Q) - 1728)^3}{j(\tau_K)^2(j(\tau_K) - 1728)^3} \right| < 877383e^{-5\pi\sqrt{|d_K|}/2}.$$

$$(iii) \text{ If } s \in \{0, 1/3, 2/3\}, \text{ then } \left| \frac{g_{\left[\begin{smallmatrix} 0 \\ 1/3 \end{smallmatrix}\right]}(\tau_K)g_{\left[\begin{smallmatrix} 1/3 \\ s \end{smallmatrix}\right]}(\tau_K)}{g_{\gamma\left[\begin{smallmatrix} 0 \\ 1/3 \end{smallmatrix}\right]}(\tau_Q)g_{\gamma\left[\begin{smallmatrix} 1/3 \\ s \end{smallmatrix}\right]}(\tau_Q)} \right| < 5.93e^{-\pi\sqrt{|d_K|}/36}.$$

Proof. (i) Since Q is a nonidentity, we have $2 \leq a \leq \sqrt{|d_K|/3}$ by Remark 4.6(ii). We then establish that

$$\begin{aligned} & \left| g_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix}\right]}(\tau_Q) \right| \\ & \geq 2t^{(1/2)\mathbf{B}_2(0)} \prod_{n=1}^{\infty} (1-t^n)^2 \quad \text{by the definition (5), where } t = e^{-\pi\sqrt{|d_K|}/a} \\ & \geq 2e^{-\pi\sqrt{|d_K|}/24} \prod_{n=1}^{\infty} (1-e^{-\pi\sqrt{3}n})^2 \quad \text{because } e^{-\pi\sqrt{|d_K|}/2} \leq t \leq e^{-\pi\sqrt{3}} \\ & \geq 2e^{-\pi\sqrt{|d_K|}/24} \prod_{n=1}^{\infty} (1+e^{-\pi 99\sqrt{3}n/100})^{-2} \\ & \quad \text{since } 1-x > 1/(1+x^{99/100}) \text{ for } 0 < x \leq e^{-\pi\sqrt{3}} \\ & \geq 2e^{-\pi\sqrt{|d_K|}/24} \prod_{n=1}^{\infty} e^{-2e^{-\pi 99\sqrt{3}n/100}} \quad \text{owing to the fact } 1+x < e^x \text{ for all } x > 0 \\ & = 2e^{-\pi\sqrt{|d_K|}/24} e^{-2e^{-\pi 99\sqrt{3}/100}/(1-e^{-\pi 99\sqrt{3}/100})} \\ & > 1.98e^{-\pi\sqrt{|d_K|}/24}. \end{aligned}$$

(ii) For convenience, let $g(\tau) = g_{\left[\begin{smallmatrix} 0 \\ 1/2 \end{smallmatrix}\right]}(\tau)^{12}$. By Lemma 5.2(i) and the fact $d_K \leq -20$, we see that

$$(14) \quad |g(\tau_K)| \leq 2^{12}e^{-\pi\sqrt{|d_K|}}e^{-24+24/(1-e^{-\pi\sqrt{20}})} < 4097e^{-\pi\sqrt{|d_K|}},$$

and similarly

$$(15) \quad \begin{aligned} |g(\tau_Q)| & \leq 2^{12}e^{-\pi\sqrt{|d_K|}/a}e^{-24+24/(1-e^{-\pi\sqrt{|d_K|}/a})} \\ & < 2^{12}e^{-\pi\sqrt{3}}e^{-24+24/(1-e^{-\pi\sqrt{3}})} < 19.71 \end{aligned}$$

by Remark 4.6(ii). It then follows from (14) and the fact $d_K \leq -20$ that

$$(16) \quad |g(\tau_K)| < 0.0033.$$

Thus we achieve that

$$\begin{aligned} & \left| \frac{j(\tau_Q)^2(j(\tau_Q) - 1728)^3}{j(\tau_K)^2(j(\tau_K) - 1728)^3} \right| \\ & = \left| \frac{g(\tau_K)^5}{g(\tau_Q)^5} \right| \left| \frac{(g(\tau_Q) + 16)^6(g(\tau_Q) + 64)^3(g(\tau_Q) - 8)^6}{(g(\tau_K) + 16)^6(g(\tau_K) + 64)^3(g(\tau_K) - 8)^6} \right| \end{aligned}$$

by Remark 3.5

$$\begin{aligned} &\leq \left(\frac{4097e^{-\pi\sqrt{|d_K|}}}{1.9812e^{-\pi\sqrt{|d_K|}/2}} \right)^5 \left| \frac{(16 + 19.71)^6(64 + 19.71)^3(8 + 19.71)^6}{(16 - 0.0033)^6(64 - 0.0033)^3(8 - 0.0033)^6} \right| \\ &\quad \text{by (i) and (14)–(16)} \\ &< 877383e^{-5\pi\sqrt{|d_K|}/2}. \end{aligned}$$

(iii) We get that

$$\begin{aligned} &\left| \frac{g_{\left[\frac{0}{1/3}\right]}(\tau_K)g_{\left[\frac{1/3}{s}\right]}(\tau_K)}{g_{\gamma\left[\frac{0}{1/3}\right]}(\tau_Q)g_{\gamma\left[\frac{1/3}{s}\right]}(\tau_Q)} \right| \\ &= \sqrt{3} \left| g_{\left[\frac{0}{1/3}\right]}(\tau_K)g_{\left[\frac{1/3}{s}\right]}(\tau_K)g_{\gamma\left[\frac{1/3}{s'}\right]}(\tau_Q)g_{\gamma\left[\frac{1/3}{s''}\right]}(\tau_Q) \right| \\ &\quad \text{with } \{s, s', s''\} = \{0, 1/3, 2/3\} \text{ by Lemma 3.11} \\ &< \sqrt{3} \left(1.75e^{-\pi\sqrt{|d_K|}/12} \right) \left(1.25e^{\pi\sqrt{|d_K|}/36} \right) \\ &\quad \times \left(1.25e^{\pi\sqrt{|d_K|}/36} \right) \left(1.25e^{\pi\sqrt{|d_K|}/36} \right) \\ &\quad \text{by Lemma 6.2 and the fact } 2 \leq a \leq \sqrt{|d_K|}/3 \\ &< 5.93e^{-\pi\sqrt{|d_K|}/36}. \quad \square \end{aligned}$$

Remark 6.4. In particular, suppose that $d_K = -20, -23$ or -24 . In this case, we have

$$C(d_K) = \begin{cases} \{x^2 + 5y^2, 2x^2 + 2xy + 3y^2\} & \text{if } d_K = -20, \\ \{x^2 + xy + 6y^2, 2x^2 \pm xy + 3y^2\} & \text{if } d_K = -23, \\ \{x^2 + 6y^2, 2x^2 + 3y^2\} & \text{if } d_K \equiv -24. \end{cases}$$

Thus, if $Q = ax^2 + bxy + cy^2$ is a nonidentity reduced form of discriminant d_K , then we obtain by Lemma 5.2(i) that

$$\begin{aligned} |g(\tau_Q)| &\leq 2^{12}e^{-\pi\sqrt{|d_K|}/a}e^{-24+24/(1-e^{-\pi\sqrt{|d_K|}/a})} \\ &\leq 2^{12}e^{-\pi\sqrt{20}/2}e^{-24+24/(1-e^{-\pi\sqrt{20}/2})} < 3.73. \end{aligned}$$

In like manner as Lemma 6.3(ii) one can derive the inequality

$$\left| \frac{j(\tau_Q)^2(j(\tau_Q) - 1728)^3}{j(\tau_K)^2(j(\tau_K) - 1728)^3} \right| < 77e^{-5\pi\sqrt{|d_K|}/2}.$$

Theorem 6.5. *If 3 splits in K , then $h_E(\varphi_E(1/3))$ generates $K_{\mathfrak{m}}$ over K .*

Proof. Since 3 splits in K , we have $d_K \equiv 1 \pmod{3}$. If $h_K = 1$, then the assertion is immediate by Proposition 4.2(iii). So, let $h_K \geq 2$, and hence

$d_K \leq -20$ by Remark 4.6(iii). By (13) and Proposition 2.3(iii) we know that there are two Galois conjugates of $f_{\begin{bmatrix} 0 \\ 1/3 \end{bmatrix}}(\tau_K)$ over H_K , namely,

$$\xi_1 = f_{\begin{bmatrix} 0 \\ 1/3 \end{bmatrix}}(\tau_K) \quad \text{and} \quad \xi_2 = \begin{cases} f_{\begin{bmatrix} 1/3 \\ 0 \end{bmatrix}}(\tau_K) & \text{if } d_K \equiv 4 \pmod{12}, \\ f_{\begin{bmatrix} 1/3 \\ 2/3 \end{bmatrix}}(\tau_K) & \text{if } d_K \equiv 1 \pmod{12}. \end{cases}$$

Let $\xi = (\xi_1 - \xi_2)^{12}$. Since ξ is fixed by every element of $\text{Gal}(K_m/H_K)$, it lies in H_K . We then derive that

$$\xi = \frac{j(\tau_K)^4(j(\tau_K) - 1728)^6}{2^{60} \cdot 3^{48}} \times \begin{cases} \frac{g_{\begin{bmatrix} 1/3 \\ 1/3 \end{bmatrix}}(\tau_K)^{12} g_{\begin{bmatrix} 1/3 \\ 2/3 \end{bmatrix}}(\tau_K)^{12}}{g_{\begin{bmatrix} 0 \\ 1/3 \end{bmatrix}}(\tau_K)^{24} g_{\begin{bmatrix} 1/3 \\ 0 \end{bmatrix}}(\tau_K)^{24}} & \text{if } d_K \equiv 4 \pmod{12}, \\ \frac{g_{\begin{bmatrix} 1/3 \\ 0 \end{bmatrix}}(\tau_K)^{12} g_{\begin{bmatrix} 1/3 \\ 1/3 \end{bmatrix}}(\tau_K)^{12}}{g_{\begin{bmatrix} 0 \\ 1/3 \end{bmatrix}}(\tau_K)^{24} g_{\begin{bmatrix} 1/3 \\ 2/3 \end{bmatrix}}(\tau_K)^{24}} & \text{if } d_K \equiv 1 \pmod{12}, \end{cases}$$

by Propositions 2.3(i) and 2.4(ii)

$$= \frac{j(\tau_K)^4(j(\tau_K) - 1728)^6}{2^{60} \cdot 3^{42}} \times \begin{cases} \frac{1}{g_{\begin{bmatrix} 0 \\ 1/3 \end{bmatrix}}(\tau_K)^{36} g_{\begin{bmatrix} 1/3 \\ 0 \end{bmatrix}}(\tau_K)^{36}} & \text{if } d_K \equiv 4 \pmod{12}, \\ \frac{1}{g_{\begin{bmatrix} 0 \\ 1/3 \end{bmatrix}}(\tau_K)^{36} g_{\begin{bmatrix} 1/3 \\ 2/3 \end{bmatrix}}(\tau_K)^{36}} & \text{if } d_K \equiv 1 \pmod{12}, \end{cases}$$

by Lemma 3.11.

Suppose that ξ does not generate H_K over K . Then, it follows from Proposition 4.5 that there is a nonidentity reduced form Q of discriminant d_K which leaves ξ fixed. Hence, we obtain that

$$1 = \left| \frac{\xi^{Q^{-1}}}{\xi} \right| = \left| \frac{j(\tau_Q)^4(j(\tau_Q) - 1728)^6}{j(\tau_K)^4(j(\tau_K) - 1728)^6} \right| \times \begin{cases} \left| \frac{g_{\begin{bmatrix} 0 \\ 1/3 \end{bmatrix}}(\tau_K)^{36} g_{\begin{bmatrix} 1/3 \\ 0 \end{bmatrix}}(\tau_K)^{36}}{g_{\gamma_Q^T \begin{bmatrix} 0 \\ 1/3 \end{bmatrix}}(\tau_Q)^{36} g_{\gamma_Q^T \begin{bmatrix} 1/3 \\ 0 \end{bmatrix}}(\tau_Q)^{36}} \right| & \text{if } d_K \equiv 4 \pmod{12}, \\ \left| \frac{g_{\begin{bmatrix} 0 \\ 1/3 \end{bmatrix}}(\tau_K)^{36} g_{\begin{bmatrix} 1/3 \\ 2/3 \end{bmatrix}}(\tau_K)^{36}}{g_{\gamma_Q^T \begin{bmatrix} 0 \\ 1/3 \end{bmatrix}}(\tau_Q)^{36} g_{\gamma_Q^T \begin{bmatrix} 1/3 \\ 2/3 \end{bmatrix}}(\tau_Q)^{36}} \right| & \text{if } d_K \equiv 1 \pmod{12}, \end{cases}$$

by Propositions 2.3(iii) and 4.5

$$< \begin{cases} \left(877383e^{-5\pi\sqrt{|d_K|/2}}\right)^2 \left(5.93e^{-\pi\sqrt{|d_K|/36}}\right)^{36} & \text{if } d_K \leq -35, \\ \left(77e^{-5\pi\sqrt{|d_K|/2}}\right)^2 \left(5.93e^{-\pi\sqrt{|d_K|/36}}\right)^{36} & \text{if } d_K = -20 \text{ or } -23, \end{cases}$$

by Lemma 6.3(ii), (iii) and Remark 6.4

< 1 ,

which yields a contradiction. Therefore, ξ generates H_K over K ; and hence $h_E(\varphi_E(1/3)) = f_{\left[\begin{smallmatrix} 0 & \\ & 1/3 \end{smallmatrix}\right]}(\tau_K)$ generates $K_{\mathfrak{m}}$ over K by Proposition 4.2(i) and (iii). \square

7. Generation of ray class fields modulo 4

Let $\mathfrak{m} = 4\mathcal{O}_K$. As a corollary of Theorems 5.1 and 5.3 we obtain the following.

Corollary 7.1. *The value $h_E(\varphi_E(1/4))$ generates $K_{\mathfrak{m}}$ over K .*

Proof. Let

$$F = K(h_E(\varphi_E(1/4))) = K\left(f_{\left[\begin{smallmatrix} 0 & \\ & 1/4 \end{smallmatrix}\right]}(\tau_K)\right)$$

which is a subfield of $K_{\mathfrak{m}}$ by Proposition 4.2(ii). Let $\min(\tau_K, \mathbb{Q}) = x^2 + bx + c$. Regarding $\gamma = \begin{bmatrix} 1-2b & -2c \\ 2 & 1 \end{bmatrix}$ as an element of $W_{K,4}/\{\pm I_2\}$ we see that

$$\begin{aligned} F &\ni \frac{1}{2} \left(f_{\left[\begin{smallmatrix} 0 & \\ & 1/4 \end{smallmatrix}\right]}(\tau_K) + f_{\left[\begin{smallmatrix} 0 & \\ & 1/4 \end{smallmatrix}\right]}(\tau_K)^\gamma \right) && \text{since } F \text{ is abelian over } K \\ &= \frac{1}{2} \left(f_{\left[\begin{smallmatrix} 0 & \\ & 1/4 \end{smallmatrix}\right]}(\tau_K) + f_{\left[\begin{smallmatrix} 2/4 & \\ & 1/4 \end{smallmatrix}\right]}(\tau_K) \right) && \text{by Propositions 4.4 and 2.3(iii)} \\ &= f_{\left[\begin{smallmatrix} 0 & \\ & 1/2 \end{smallmatrix}\right]}(\tau_K) && \text{by Lemma 3.8.} \end{aligned}$$

Thus we achieve that

$$\begin{aligned} K_{\mathfrak{m}} &\supseteq F \\ &= K\left(f_{\left[\begin{smallmatrix} 0 & \\ & 1/4 \end{smallmatrix}\right]}(\tau_K), f_{\left[\begin{smallmatrix} 0 & \\ & 1/2 \end{smallmatrix}\right]}(\tau_K)\right) \\ &= K_{2\mathcal{O}_K}\left(f_{\left[\begin{smallmatrix} 0 & \\ & 1/4 \end{smallmatrix}\right]}(\tau_K)\right) && \text{by Theorems 5.1 and 5.3} \\ &= K_{\mathfrak{m}} && \text{by Proposition 4.2(iii),} \end{aligned}$$

and hence $F = K_{\mathfrak{m}}$. This completes the proof. \square

8. Generation of ray class fields modulo 6

Let $\mathfrak{m} = 6\mathcal{O}_K$. As corollaries of Theorems 5.1, 5.3, 6.1 and 6.5 we obtain the followings.

Corollary 8.1. *If 2 splits in K , then $h_E(\varphi_E(1/3))$ generates $K_{\mathfrak{m}}$ over K .*

Proof. We see that

$$\begin{aligned} [K_{\mathfrak{m}} : K_{3\mathcal{O}_K}] &= \frac{[K_{\mathfrak{m}} : H_K]}{[K_{3\mathcal{O}_K} : H_K]} \\ &= \phi(2\mathcal{O}_K) \quad \text{by Lemma 4.1 and the fact } \mathcal{O}_K^* = \{\pm 1\} \\ &= 1 \quad \text{because we are assuming that 2 splits in } K, \end{aligned}$$

from which it follows that $K_{\mathfrak{m}} = K_{3\mathcal{O}_K}$. Thus, we conclude by Theorems 6.1 and 6.5 that

$$K_{\mathfrak{m}} = K_{3\mathcal{O}_K} = K(h_E(\varphi_E(1/3))). \quad \square$$

Corollary 8.2. *Suppose that 2 is ramified and 3 is inert in K . Then $h_E(\varphi_E(1/6))$ generates $K_{\mathfrak{m}}$ over K .*

Proof. Let

$$F = K(h_E(\varphi_E(1/6))) = K\left(f_{\begin{bmatrix} 0 \\ 1/6 \end{bmatrix}}(\tau_K)\right)$$

which is a subfield of $K_{\mathfrak{m}}$ by Proposition 4.2(ii). Now that $d_K \equiv 8 \pmod{12}$, one can consider the matrices

$$\alpha = \begin{bmatrix} 1 & d_K/2 \\ 2 & 1 \end{bmatrix}, \quad \beta = \begin{bmatrix} 3 & d_K/2 \\ 2 & 3 \end{bmatrix}, \quad \gamma = \begin{bmatrix} 5 & d_K/2 \\ 2 & 5 \end{bmatrix}$$

as elements of $W_{K,6}/\{\pm I_2\}$. Since F is an abelian extension of K , we obtain that

$$\begin{aligned} F &\ni \frac{1}{4} \left(f_{\begin{bmatrix} 0 \\ 1/6 \end{bmatrix}}(\tau_K) + f_{\begin{bmatrix} 0 \\ 1/6 \end{bmatrix}}(\tau_K)^\alpha + f_{\begin{bmatrix} 0 \\ 1/6 \end{bmatrix}}(\tau_K)^\beta + f_{\begin{bmatrix} 0 \\ 1/6 \end{bmatrix}}(\tau_K)^\gamma \right) \\ &= \frac{1}{4} \left(f_{\begin{bmatrix} 0 \\ 1/6 \end{bmatrix}}(\tau_K) + f_{\begin{bmatrix} 2/6 \\ 1/6 \end{bmatrix}}(\tau_K) + f_{\begin{bmatrix} 2/6 \\ 3/6 \end{bmatrix}}(\tau_K) + f_{\begin{bmatrix} 2/6 \\ 5/6 \end{bmatrix}}(\tau_K) \right) \\ &\quad \text{by Propositions 4.4 and 2.3(iii)} \\ &= f_{\begin{bmatrix} 0 \\ 1/2 \end{bmatrix}}(\tau_K) \quad \text{by Lemma 3.9.} \end{aligned}$$

We then attain that

$$\begin{aligned} K_{\mathfrak{m}} &\supseteq F \\ &= K\left(f_{\begin{bmatrix} 0 \\ 1/6 \end{bmatrix}}(\tau_K), f_{\begin{bmatrix} 0 \\ 1/2 \end{bmatrix}}(\tau_K)\right) \\ &= K_{2\mathcal{O}_K}\left(f_{\begin{bmatrix} 0 \\ 1/6 \end{bmatrix}}(\tau_K)\right) \quad \text{by Theorems 5.1 and 5.3} \\ &= K_{\mathfrak{m}} \quad \text{by Proposition 4.2(iii),} \end{aligned}$$

from which it follows that $F = K_{\mathfrak{m}}$. \square

Corollary 8.3. *If 2 is inert in K , then $h_E(\varphi_E(1/6))$ generates $K_{\mathfrak{m}}$ over K .*

Proof. Let

$$F = K(h_E(\varphi_E(1/6))) = K\left(f_{\left[\begin{smallmatrix} 0 \\ 1/6 \end{smallmatrix}\right]}(\tau_K)\right)$$

which is a subfield of K_m by Proposition 4.2(ii). Since 2 is inert in K , we have $d_K \equiv 5 \pmod{8}$ and $\tau_K = (-1 + \sqrt{d_K})/2$. Let

$$\alpha = \begin{bmatrix} 4 & 3(d_K - 1)/4 \\ 3 & 1 \end{bmatrix} \quad \text{and} \quad \beta = \begin{bmatrix} 1 & 3(d_K - 1)/4 \\ 3 & 4 \end{bmatrix}$$

which are elements of $W_{K,6}/\{\pm I_2\}$. Since F is an abelian extension of K , we get that

$$\begin{aligned} F &\ni \frac{1}{3} \left(f_{\left[\begin{smallmatrix} 0 \\ 1/6 \end{smallmatrix}\right]}(\tau_K) + f_{\left[\begin{smallmatrix} 0 \\ 1/6 \end{smallmatrix}\right]}(\tau_K)^\alpha + f_{\left[\begin{smallmatrix} 0 \\ 1/6 \end{smallmatrix}\right]}(\tau_K)^\beta \right) \\ &= \frac{1}{3} \left(f_{\left[\begin{smallmatrix} 0 \\ 1/6 \end{smallmatrix}\right]}(\tau_K) + f_{\left[\begin{smallmatrix} 3/6 \\ 1/6 \end{smallmatrix}\right]}(\tau_K) + f_{\left[\begin{smallmatrix} 3/6 \\ 4/6 \end{smallmatrix}\right]}(\tau_K) \right) \\ &\quad \text{by Propositions 4.4 and 2.3(iii)} \\ &= f_{\left[\begin{smallmatrix} 0 \\ 1/3 \end{smallmatrix}\right]}(\tau_K) \quad \text{by Lemma 3.10.} \end{aligned}$$

Whence we obtain that

$$\begin{aligned} K_m &\supseteq F \\ &= K\left(f_{\left[\begin{smallmatrix} 0 \\ 1/6 \end{smallmatrix}\right]}(\tau_K), f_{\left[\begin{smallmatrix} 0 \\ 1/3 \end{smallmatrix}\right]}(\tau_K)\right) \\ &= K_{3\mathcal{O}_K}\left(f_{\left[\begin{smallmatrix} 0 \\ 1/6 \end{smallmatrix}\right]}(\tau_K)\right) \quad \text{by Theorems 6.1 and 6.5} \\ &= K_m \quad \text{by Proposition 4.2(iii).} \end{aligned}$$

Therefore, we achieve $F = K_m$. \square

Lemma 8.4. *Suppose that $d_K \leq -20$ and $h_K \geq 2$. Let $Q = ax^2 + bxy + cy^2$ be a nonidentity reduced form of discriminant d_K , and let $\begin{bmatrix} r \\ s \end{bmatrix} \in (1/6)\mathbb{Z}^2 \setminus \mathbb{Z}^2$.*

(i) *If $\langle r \rangle \neq 0$ and $t = |e^{2\pi i \tau_K}|$, then*

$$\left| g_{\begin{bmatrix} r \\ s \end{bmatrix}}(\tau_K) \right| < 1.1t^{(1/2)\mathbf{B}_2(\langle r \rangle)}.$$

(ii) *If $t_Q = |e^{2\pi i \tau_Q}|$, then*

$$\left| g_{\begin{bmatrix} r \\ s \end{bmatrix}}(\tau_Q) \right| > 0.75t_Q^{1/12}.$$

(iii) *Furthermore, we have*

$$\left| \frac{g_{\left[\begin{smallmatrix} 0 \\ 1/6 \end{smallmatrix}\right]}(\tau_K)}{g_{\begin{bmatrix} r \\ s \end{bmatrix}}(\tau_Q)} \right| < 1.$$

Proof. (i) We deduce that

$$\begin{aligned}
|g_{[r_s]}(\tau_K)| &\leq t^{(1/2)\mathbf{B}_2(\langle r \rangle)}(1+t^{1/6}) \prod_{n=1}^{\infty} (1+t^{n-1/2})^2 \\
&\quad \text{by the definition (5) and Proposition 2.3(i)} \\
&\leq t^{(1/2)\mathbf{B}_2(\langle r \rangle)}(1+t^{1/6}) \prod_{n=1}^{\infty} e^{2t^{n-1/2}} \quad \text{since } 1+x < e^x \text{ for all } x > 0 \\
&= t^{(1/2)\mathbf{B}_2(\langle r \rangle)}(1+t^{1/6})e^{2t^{1/2}/(1-t)} \\
&< 1.1t^{(1/2)\mathbf{B}_2(\langle r \rangle)} \quad \text{because } t = e^{-\pi\sqrt{|d_K|}} \leq e^{-\pi\sqrt{20}}.
\end{aligned}$$

(ii) We further derive that

$$\begin{aligned}
|g_{[r_s]}(\tau_Q)| &\geq \begin{cases} |1-\zeta_6|t_Q^{(1/2)\mathbf{B}_2(0)} \prod_{n=1}^{\infty} (1-t_Q^n)^2 & \text{if } \langle r \rangle = 0, \\ t_Q^{(1/2)\mathbf{B}_2(\langle r \rangle)}(1-t_Q^{1/6}) \prod_{n=1}^{\infty} (1-t_Q^{n-1/2})^2 & \text{if } \langle r \rangle \neq 0, \end{cases} \\
&\quad \text{by the definition (5) and Proposition 2.3(i)} \\
&\geq \begin{cases} t_Q^{1/12} \prod_{n=1}^{\infty} (1-e^{-\pi\sqrt{3}n})^2 & \text{if } \langle r \rangle = 0, \\ t_Q^{(1/2)\mathbf{B}_2(\langle r \rangle)}(1-e^{-\pi\sqrt{3}/6})(1-e^{-\pi\sqrt{3}/2})^2 \\ \quad \times \prod_{n=2}^{\infty} (1-e^{-\pi\sqrt{3}(n-1/2)})^2 & \text{if } \langle r \rangle \neq 0, \end{cases} \\
&\quad \text{because } t_Q = e^{-\pi\sqrt{|d_K|}/a} \leq e^{-\pi\sqrt{3}} \text{ by Remark 4.6(ii)} \\
&\geq \begin{cases} t_Q^{1/12} \prod_{n=1}^{\infty} (1+e^{-\pi 99\sqrt{3}n/100})^{-2} & \text{if } \langle r \rangle = 0, \\ t_Q^{(1/2)\mathbf{B}_2(\langle r \rangle)}(1-e^{-\pi\sqrt{3}/6})(1-e^{-\pi\sqrt{3}/2})^2 \\ \quad \times \prod_{n=2}^{\infty} (1+e^{-\pi 99\sqrt{3}(n-1/2)/100})^{-2} & \text{if } \langle r \rangle \neq 0, \end{cases} \\
&\quad \text{by the fact } 1-x > 1/(1+x^{99/100}) \text{ for } 0 < x \leq e^{-\pi\sqrt{3}} \\
&\geq \begin{cases} t_Q^{1/12} \prod_{n=1}^{\infty} e^{-2e^{-\pi 99\sqrt{3}n/100}} & \text{if } \langle r \rangle = 0, \\ t_Q^{(1/2)\mathbf{B}_2(\langle r \rangle)}(1-e^{-\pi\sqrt{3}/6})(1-e^{-\pi\sqrt{3}/2})^2 \\ \quad \times \prod_{n=2}^{\infty} e^{-2e^{-\pi 99\sqrt{3}(n-1/2)/100}} & \text{if } \langle r \rangle \neq 0, \end{cases} \\
&\quad \text{since } 1+x < e^x \text{ for all } x > 0
\end{aligned}$$

$$\begin{aligned}
&= \begin{cases} t_Q^{1/12} e^{-2e^{-\pi 99\sqrt{3}/100}/(1-e^{-\pi 99\sqrt{3}/100})} & \text{if } \langle r \rangle = 0, \\ t_Q^{(1/2)\mathbf{B}_2(\langle r \rangle)} (1 - e^{-\pi\sqrt{3}/6})(1 - e^{-\pi\sqrt{3}/2})^2 \\ \times e^{-2e^{-\pi 297\sqrt{3}/200}/(1-e^{-\pi 99\sqrt{3}/100})} & \text{if } \langle r \rangle \neq 0 \end{cases} \\
&> \begin{cases} 0.99t_Q^{1/12} & \text{if } \langle r \rangle = 0, \\ 0.519t_Q^{(1/2)\mathbf{B}_2(\langle r \rangle)} & \text{if } \langle r \rangle \neq 0 \end{cases} \\
&\geq \begin{cases} 0.99t_Q^{1/12} & \text{if } \langle r \rangle = 0, \\ 0.519t_Q^{1/72} & \text{if } \langle r \rangle \neq 0 \end{cases} \\
&> 0.75t_Q^{1/12} \quad \text{due to the fact } t_Q \leq e^{-\pi\sqrt{3}}.
\end{aligned}$$

(iii) See [8, Lemma 4.1]. \square

Theorem 8.5. *If 2 is ramified in K and 3 is not inert in K , then $h_E(\varphi_E(1/6))$ generates K_m over K .*

Proof. If $h_K = 1$, then the assertion is straightforward by Proposition 4.2(iii). Now, let $h_K \geq 2$. Since 2 is ramified in K and 3 is not inert in K , we have $d_K \equiv 0$ or $4 \pmod{12}$, and hence $d_K \leq -20$ by Remark 4.6(iii). Let

$$F = K(h_E(\varphi_E(1/6))) = K\left(f_{\begin{bmatrix} 0 \\ 1/6 \end{bmatrix}}(\tau_K)\right)$$

which is an abelian extension of K as a subfield of K_m by Proposition 4.2(ii). Let α be the element of $W_{K,6}/\{\pm I_2\}$ given by

$$\alpha = \begin{cases} \begin{bmatrix} 1 & d_K \\ 4 & 1 \end{bmatrix} & \text{if } d_K \equiv 0 \pmod{12}, \\ \begin{bmatrix} 3 & d_K/2 \\ 2 & 3 \end{bmatrix} & \text{if } d_K \equiv 4 \pmod{12}, \end{cases}$$

and set

$$\xi = \left(f_{\begin{bmatrix} 0 \\ 1/6 \end{bmatrix}}(\tau_K)^\alpha - f_{\begin{bmatrix} 0 \\ 1/6 \end{bmatrix}}(\tau_K)\right)^{12}$$

which belongs to F . We then derive that

$$\begin{aligned}
\xi &= \begin{cases} \left(f_{\begin{bmatrix} 4/6 \\ 1/6 \end{bmatrix}}(\tau_K) - f_{\begin{bmatrix} 0 \\ 1/6 \end{bmatrix}}(\tau_K)\right)^{12} & \text{if } d_K \equiv 0 \pmod{12}, \\ \left(f_{\begin{bmatrix} 2/6 \\ 3/6 \end{bmatrix}}(\tau_K) - f_{\begin{bmatrix} 0 \\ 1/6 \end{bmatrix}}(\tau_K)\right)^{12} & \text{if } d_K \equiv 4 \pmod{12}, \end{cases} \\
&\quad \text{by Propositions 4.4 and 2.3(iii)} \\
&= \frac{j(\tau_K)^4(j(\tau_K) - 1728)^6}{2^{60} \cdot 3^{48}}
\end{aligned}$$

$$\begin{aligned}
& \times \begin{cases} \frac{g_{\left[\frac{1/3}{2/3}\right]}(\tau_K)^{12} g_{\left[\frac{1/3}{0}\right]}(\tau_K)^{12}}{g_{\left[\frac{1/3}{5/6}\right]}(\tau_K)^{24} g_{\left[\frac{0}{1/6}\right]}(\tau_K)^{24}} & \text{if } d_K \equiv 0 \pmod{12}, \\ \frac{g_{\left[\frac{1/3}{2/3}\right]}(\tau_K)^{12} g_{\left[\frac{1/3}{1/3}\right]}(\tau_K)^{12}}{g_{\left[\frac{1/3}{1/2}\right]}(\tau_K)^{24} g_{\left[\frac{0}{1/6}\right]}(\tau_K)^{24}} & \text{if } d_K \equiv 4 \pmod{12}, \end{cases} \\
& \text{by Proposition 2.4(ii)} \\
& = \frac{j(\tau_K)^4(j(\tau_K) - 1728)^6}{2^{60} \cdot 3^{48}} \\
& \times \begin{cases} \frac{3^6}{g_{\left[\frac{0}{1/3}\right]}(\tau_K)^{12} g_{\left[\frac{1/3}{1/3}\right]}(\tau_K)^{12} g_{\left[\frac{1/3}{5/6}\right]}(\tau_K)^{24} g_{\left[\frac{0}{1/6}\right]}(\tau_K)^{24}} & \text{if } d_K \equiv 0 \pmod{12}, \\ \frac{3^6}{g_{\left[\frac{0}{1/3}\right]}(\tau_K)^{12} g_{\left[\frac{1/3}{0}\right]}(\tau_K)^{12} g_{\left[\frac{1/3}{1/2}\right]}(\tau_K)^{24} g_{\left[\frac{0}{1/6}\right]}(\tau_K)^{24}} & \text{if } d_K \equiv 4 \pmod{12}, \end{cases} \\
& \text{by Lemma 3.11.}
\end{aligned}$$

Suppose on the contrary that $F = K\left(f_{\left[\frac{0}{1/6}\right]}(\tau_K)\right) = K\left(f_{\left[\frac{0}{1/6}\right]}(\tau_K), \xi\right)$ is properly contained in K_m . Then, by Propositions 4.4 and 4.5 there is a pair $(\gamma, Q) \in W_{K,6}/\{\pm I_2\} \times C(d_K)$ for which γQ^{-1} leaves both $f_{\left[\frac{0}{1/6}\right]}(\tau_K)$ and ξ fixed. Observe that Proposition 4.2(iii) implies that $Q = ax^2 + bxy + cy^2$ must be a nonidentity reduced form. Let

$$t = |\tau_K| = e^{-\pi\sqrt{|d_K|}} \quad \text{and} \quad t_Q = |\tau_Q| = e^{-\pi\sqrt{|d_K|}/a}.$$

Then we achieve that

$$\begin{aligned}
1 &= \left| \frac{\xi^{\gamma Q^{-1}}}{\xi} \right| = \left| \frac{j(\tau_Q)^4(j(\tau_Q) - 1728)^6}{j(\tau_K)^4(j(\tau_K) - 1728)^6} \right| \left| \frac{g_{\left[\frac{0}{1/6}\right]}(\tau_K)^{24}}{g_{\gamma_Q^T \gamma^T \left[\frac{0}{1/6}\right]}(\tau_Q)^{24}} \right| \\
& \times \begin{cases} \left| \frac{g_{\left[\frac{0}{1/3}\right]}(\tau_K)^{12} g_{\left[\frac{1/3}{1/3}\right]}(\tau_K)^{12}}{g_{\gamma_Q^T \gamma^T \left[\frac{0}{1/3}\right]}(\tau_Q)^{12} g_{\gamma_Q^T \gamma^T \left[\frac{1/3}{1/3}\right]}(\tau_Q)^{12}} \right| \left| \frac{g_{\left[\frac{1/3}{5/6}\right]}(\tau_K)^{24}}{g_{\gamma_Q^T \gamma^T \left[\frac{1/3}{5/6}\right]}(\tau_Q)^{24}} \right| & \text{if } d_K \equiv 0 \pmod{12}, \\ \left| \frac{g_{\left[\frac{0}{1/3}\right]}(\tau_K)^{12} g_{\left[\frac{1/3}{0}\right]}(\tau_K)^{12}}{g_{\gamma_Q^T \gamma^T \left[\frac{0}{1/3}\right]}(\tau_Q)^{12} g_{\gamma_Q^T \gamma^T \left[\frac{1/3}{0}\right]}(\tau_Q)^{12}} \right| \left| \frac{g_{\left[\frac{1/3}{1/2}\right]}(\tau_K)^{24}}{g_{\gamma_Q^T \gamma^T \left[\frac{1/3}{1/2}\right]}(\tau_Q)^{24}} \right| & \text{if } d_K \equiv 4 \pmod{12}, \end{cases}
\end{aligned}$$

by Propositions 4.4 and 2.3(iii)

$$< \begin{cases} \left(77e^{-5\pi\sqrt{|d_K|/2}}\right)^2 \\ \times \left(5.93e^{-\pi\sqrt{|d_K|/36}}\right)^{12} \left(1.1t^{(1/2)\mathbf{B}_2(1/3)}\right)^{24} \left(0.75t_Q^{1/12}\right)^{-24} \\ \text{if } d_K = -20 \text{ or } -24, \\ \left(877383e^{-5\pi\sqrt{|d_K|/2}}\right)^2 \\ \times \left(5.93e^{-\pi\sqrt{|d_K|/36}}\right)^{12} \left(1.1t^{(1/2)\mathbf{B}_2(1/3)}\right)^{24} \left(0.75t_Q^{1/12}\right)^{-24} \\ \text{if } d_K \leq -56, \end{cases}$$

by Lemma 6.3(ii), (iii), Remarks 6.4 and Lemma 8.4

$$\begin{aligned} &< e^{-(14/3-2/a)\pi\sqrt{|d_K|}} \times \begin{cases} 1.1006 \cdot 10^{17} & \text{if } d_K = -20 \text{ or } -24, \\ 1.429 \cdot 10^{25} & \text{if } d_K \leq -56 \end{cases} \\ &\leq e^{-(11/3)\pi\sqrt{|d_K|}} \times \begin{cases} 1.1006 \cdot 10^{17} & \text{if } d_K = -20 \text{ or } -24, \\ 1.429 \cdot 10^{25} & \text{if } d_K \leq -56, \end{cases} \quad \text{since } a \geq 2 \\ &< 1, \end{aligned}$$

which yields a contradiction. Therefore, we conclude that $F = K_{\mathfrak{m}}$. \square

References

- [1] A. Baker, *Linear forms in the logarithms of algebraic numbers I*, *Mathematika* **13** (1966), 204–216.
- [2] D. A. Cox, *Primes of the form $x^2 + ny^2$* , A Wiley-Interscience Publication, John Wiley & Sons, Inc., New York, 1989.
- [3] M. Deuring, *Die Klassenkörper der komplexen Multiplikation*, *Enzyklopädie der mathematischen Wissenschaften: Mit Einschluss ihrer Anwendungen*, Band I 2, Heft 10, Teil II (Article I 2, 23), B. G. Teubner Verlagsgesellschaft, Stuttgart, 1958.
- [4] G. Frei, F. Lemmermeyer, and P. J. Roquette, *Emil Artin and Helmut Hasse—the correspondence 1923–1958*, *Contributions in Mathematical and Computational Sciences*, 5, Springer, Heidelberg, 2014.
- [5] H. Hasse, *Neue Begründung der komplexen Multiplikation. Zweiter Teil: Aufbau ohne Benutzung der allgemeinen Klassenkörpertheorie*, *J. Reine Angew. Math.* **165** (1931), 64–88.
- [6] K. Heegner, *Diophantische Analysis und Modulfunktionen*, *Math. Z.* **56** (1952), 227–253.
- [7] G. J. Janusz, *Algebraic number fields*, second edition, *Graduate Studies in Mathematics*, 7, American Mathematical Society, Providence, RI, 1996.
- [8] H. Y. Jung, J. K. Koo, and D. H. Shin, *Normal bases of ray class fields over imaginary quadratic fields*, *Math. Z.* **271** (2012), no. 1-2, 109–116.
- [9] C. H. Kim and J. K. Koo, *On the genus of some modular curves of level N* , *Bull. Austral. Math. Soc.* **54** (1996), no. 2, 291–297.
- [10] ———, *Arithmetic of the modular function $j_{1,8}$* , *Ramanujan J.* **4** (2000), no. 3, 317–338.
- [11] J. K. Koo and D. H. Shin, *On some arithmetic properties of Siegel functions*, *Math. Z.* **264** (2010), no. 1, 137–177.
- [12] J. K. Koo, D. H. Shin, and D. S. Yoon, *Generation of class fields by using the Weber function*, *J. Number Theory* **167** (2016), 74–87.
- [13] ———, *On a problem of Hasse and Ramachandra*, submitted, <https://arxiv.org/abs/1608.06705>.

- [14] D. S. Kubert and S. Lang, *Modular units*, Grundlehren der Mathematischen Wissenschaften, 244, Springer-Verlag, New York, 1981.
- [15] S. Lang, *Elliptic functions*, second edition, Graduate Texts in Mathematics, 112, Springer-Verlag, New York, 1987.
- [16] ———, *Algebraic number theory*, second edition, Graduate Texts in Mathematics, 110, Springer-Verlag, New York, 1994.
- [17] K. Ramachandra, *Some applications of Kronecker's limit formulas*, Ann. of Math. (2) **80** (1964), 104–148.
- [18] G. Shimura, *Introduction to the arithmetic theory of automorphic functions*, Publications of the Mathematical Society of Japan, No. 11. Iwanami Shoten, Publishers, Tokyo, 1971.
- [19] J. H. Silverman, *The arithmetic of elliptic curves*, second edition, Graduate Texts in Mathematics, 106, Springer, Dordrecht, 2009.
- [20] H. M. Stark, *A complete determination of the complex quadratic fields of class-number one*, Michigan Math. J. **14** (1967), 1–27.
- [21] ———, *On the “gap” in a theorem of Heegner*, J. Number Theory **1** (1969), 16–27.
- [22] P. Stevenhagen, *Hilbert's 12th problem, complex multiplication and Shimura reciprocity*, in Class field theory—its centenary and prospect (Tokyo, 1998), 161–176, Adv. Stud. Pure Math., 30, Math. Soc. Japan, Tokyo, 2001.
- [23] M. Sugawara, *On the so-called Kronecker's dream in young days*, Proc. Phys.-Math. Japan **15** (1933), no. 3, 99–107.
- [24] ———, *Zur Theorie der komplexen Multiplikation. I*, J. Reine Angew. Math. **174** (1936), 189–191.

HO YUN JUNG
 APPLIED ALGEBRA AND OPTIMIZATION RESEARCH CENTER
 SUNGKYUNKWAN UNIVERSITY
 SUWON-SI 16419, KOREA
E-mail address: hoyunjung@skku.edu

JA KYUNG KOO
 DEPARTMENT OF MATHEMATICAL SCIENCES
 KAIST
 DAEJEON 34141, KOREA
E-mail address: jkkoo@math.kaist.ac.kr

DONG HWA SHIN
 DEPARTMENT OF MATHEMATICS
 HANKUK UNIVERSITY OF FOREIGN STUDIES
 YONGIN-SI 17035, KOREA
E-mail address: dhshin@hufs.ac.kr