

## SOME CONDITIONS ON THE FORM OF THIRD ELEMENT FROM DIOPHANTINE PAIRS AND ITS APPLICATION

JUNE BOK LEE AND JINSEO PARK

ABSTRACT. A set  $\{a_1, a_2, \dots, a_m\}$  of positive integers is called a Diophantine  $m$ -tuple if  $a_i a_j + 1$  is a perfect square for all  $1 \leq i < j \leq m$ . In this paper, we show that the form of third element in Diophantine pairs and develop some results which are needed to prove the extendibility of the Diophantine pair  $\{a, b\}$  with some conditions. By using this result, we prove the extendibility of Diophantine pairs  $\{F_{k-2}F_{k+1}, F_{k-1}F_{k+2}\}$  and  $\{F_{k-2}F_{k-1}, F_{k+1}F_{k+2}\}$ , where  $F_n$  is the  $n$ -th Fibonacci number.

### 1. Introduction

A Diophantine  $m$ -tuple is a set which consists of  $m$  distinct positive integers satisfying the property that the product of any two of them is one less than a perfect square. If the set which consists of rational numbers satisfy the same property, then we called rational Diophantine  $m$ -tuple. Diophantus found the first rational Diophantine quadruple  $\{1/16, 33/16, 17/4, 105/16\}$ . However, the first Diophantine quadruple  $\{1, 3, 8, 120\}$  was found by Fermat. Many famous mathematicians made lots of results related to the problems of Diophantine  $m$ -tuple, but still there are many open problems. Especially, the most famous problem is the extendibility of Diophantine  $m$ -tuple.

For any Diophantine triple  $\{a, b, c\}$  with  $a < b < c$ , the set  $\{a, b, c, d_{\pm}\}$  is a Diophantine quadruple, where

$$d_{\pm} = a + b + c + 2abc \pm 2rst$$

and  $r, s, t$  are the positive integers satisfying

$$ab + 1 = r^2, \quad ac + 1 = s^2, \quad bc + 1 = t^2.$$

A folklore conjecture is that there does not exist a Diophantine quintuple. The stronger version of this conjecture states that if  $\{a, b, c, d\}$  is a Diophantine

---

Received April 29, 2017; Accepted June 19, 2017.

2010 *Mathematics Subject Classification.* Primary 11B39, 11D09, 11D45; Secondary 11B37, 11J68, 11J86.

*Key words and phrases.* Diophantine  $m$ -tuple, Fibonacci numbers, Pell equation.

The second named author was also supported by the National Research Foundation of Korea(NRF) grant founded by the Korea government(NRF-2013R1A1A2063240).

quadruple and  $d > \max\{a, b, c\}$ , then  $d = d_+$ . These Diophantine quadruples are called regular.

The reason why the extendibility is important is related to the elliptic curves. To extend the Diophantine triple  $\{a, b, c\}$  to Diophantine quadruple, we have to solve the equations

$$ax + 1 = \square, \quad bx + 1 = \square, \quad cx + 1 = \square.$$

Hence, we have the equation

$$E : y^2 = (ax + 1)(bx + 1)(cx + 1),$$

which is the elliptic curve by the product of three equations. Then we have always the integer points

$$(0, \pm 1), \quad (d_+, \pm(at + rs)(bs + rt)(cr + st)), \quad (d_-, \pm((at - rs)(bs - rt)(cr - st))),$$

and also  $(-1, 0)$  if  $1 \in \{a, b, c\}$ . The conjecture means it is possible to prove that there are no other integer points on  $E$  for some family of Diophantine triples. For example, A. Dujella [5] proved that the elliptic curve

$$E_k : y^2 = ((k - 1)x + 1)((k + 1)x + 1)(4kx + 1)$$

has four integer points

$$(0, \pm 1), \quad (16k^3 - 4k, \pm(128k^6 - 112k^4 - 20k^2 - 1))$$

under assumption that  $\text{rank}(E_k(\mathbb{Q})) = 1$ . Similar results [7] and [13] were proved for the equation

$$y^2 = (F_{2k}x + 1)(F_{2k+2}x + 1)(F_{2k+4} + 1)$$

and

$$y^2 = (F_{2k+1}x + 1)(F_{2k+3}x + 1)(F_{2k+5} + 1),$$

respectively, where  $F_n$  is the  $n$ -th Fibonacci number, defined by  $F_0 = 0, F_1 = 1$  and  $F_{n+2} = F_{n+1} + F_n$ . Therefore, it is important not only how to prove the extendibility of Diophantine  $m$ -tuple but also which case of Diophantine  $m$ -tuple is proved.

In 1969, A. Baker and H. Davenport [1] proved that the Diophantine triple  $\{1, 3, 8\}$  can not be extended to a Diophantine quintuple, namely, if the set  $\{1, 3, 8, d\}$  is a Diophantine quadruple, then  $d = 120$ . This means the Diophantine triple  $\{1, 3, 8\}$  is regular.

Let us consider the Diophantine pair  $\{1, 3\}$ . If the set  $\{1, 3, c\}$  is a Diophantine triple, then we have the equation

$$y^2 - 3x^2 = -2,$$

since  $c + 1 = x^2$ ,  $3c + 1 = y^2$ . We easily find the form of solutions of the Diophantine equation above is

$$(y + x\sqrt{3}) = (y_0 + x_0\sqrt{3})(2 + \sqrt{3})^k.$$

If  $(y_0, x_0)$  belongs to the same class as either of the solutions  $(\pm 1, 1)$ , then we have the form of third element

$$c = c_k = \frac{1}{6}[(2 + \sqrt{3})(7 + 4\sqrt{3})^k + (2 - \sqrt{3})(7 - 4\sqrt{3})^k - 4].$$

In 1998, Dujella and A. Pethö [9] developed this result, that is, if the set  $\{1, 3, c_k, d\}$  is a Diophantine quadruple, then  $d = d_- = c_{k-1}$  or  $d_+ = c_{k+1}$ . This result shows how important the form of third element in Diophantine pair is. Then we have a question such that how can we find the form of third element in Diophantine pairs. The form of third element  $c_k$  in the Diophantine pair  $\{1, 3\}$  is found by using Theorem 8 in [15]. In fact, for a fixed Diophantine pair  $\{a, b\}$  with  $a < b$ , the theorem shows the form of third element, where  $b < 4a$ . This theorem was also used in [3], [4], [12].

The recent research of third element in Diophantine  $m$ -tuple is Lemma 4.1 in [10] which shows the form of third element in the Diophantine pair  $\{a, b\}$ , where  $a < b \leq 8a$ . Unfortunately however, the general form of third elements are unknown until now.

In this paper, we use the following identity

$$F_{k-2}F_{k-1}F_{k+1}F_{k+2} + 1 = F_k^4,$$

which was proved by E. Cesàro. We can make many Diophantine pairs by using above identity. However, we consider only two sets

$$\{F_{k-2}F_{k+1}, F_{k-1}F_{k+2}\} \quad \text{and} \quad \{F_{k-2}F_{k-1}, F_{k+1}F_{k+2}\},$$

since the first set satisfy the condition of Lemma 4.1 in [10], namely,  $F_{k-1}F_{k+2} < 8F_{k-2}F_{k+1}$ , but the second set does not satisfy the condition. Hence, we need to generalize the Lemma 4.1 for the extendibility of Diophantine pairs which do not satisfy the condition  $b \leq 8a$ . Also, we develop some results which need to prove the extendibility of Diophantine pair  $\{a, b\}$  with some conditions. Through these generalized result, we prove the extendibility of  $\{F_{k-2}F_{k+1}, F_{k-1}F_{k+2}\}$  and  $\{F_{k-2}F_{k-1}, F_{k+1}F_{k+2}\}$ .

## 2. Preliminaries

### 2.1. The bounds of each elements of Diophantine triple

We can find the lower bounds of second element of the Diophantine triple  $\{a, b, c\}$  with  $a < b$  using the following lemma.

**Lemma 2.1** ([11, Lemma 1.3]). *Suppose that  $\{a, b, c, d\}$  is a Diophantine quadruple with  $a < b < c < d_+ < d$ .*

- (1) *If  $b < 2a$ , then  $b > 2.1 \cdot 10^4$ .*
- (2) *If  $2a \leq b \leq 8a$ , then  $b > 1.3 \cdot 10^5$ .*
- (3) *If  $b > 8a$ , then  $b > 2 \cdot 10^3$ .*

Let  $\{a, b, c\}$  be a Diophantine triple, and  $r, s, t$  be the positive integers satisfying  $ab + 1 = r^2$ ,  $ac + 1 = s^2$ ,  $bc + 1 = t^2$ . Then we have

$$at^2 - bs^2 = a - b.$$

We easily find the form of solutions of the equation above is

$$(t\sqrt{a} + s\sqrt{b}) = (t_0\sqrt{a} + s_0\sqrt{b})(r + \sqrt{bc})^\nu.$$

If  $(t_0, s_0)$  belongs to the same class as either of the solutions  $(\pm 1, 1)$ , then  $s$  can be expressed as  $s = s_\nu^\tau$ , where  $\tau \in \{\pm 1\}$  and

$$s_0 = s_0^\tau = 1, \quad s_1^\tau = r + \tau a, \quad s_{\nu+2}^\tau = 2rs_{\nu+1}^\tau - s_\nu^\tau.$$

Define  $c_\nu^\tau = ((s_\nu^\tau)^2 - 1)/a$ . Then, we obtain

$$c = c_\nu^\tau = \frac{1}{4ab} [(a + b \pm 2\sqrt{ab})(2ab + 1 + 2r\sqrt{ab})^\nu + (a + b \mp 2\sqrt{ab})(2ab + 1 - 2r\sqrt{ab})^\nu - 2(a + b)].$$

First, we have the form of third element  $c$  in the Diophantine triple  $\{a, b, c\}$  by the following theorem.

**Theorem 2.2** ([15, Theorem 8]). *If  $a < b < 4a$ , and  $b$  are in  $\mathbb{Z}^+$ , and*

$$ab + 1 = r^2, \quad ac + 1 = s^2, \quad bc + 1 = t^2$$

*holds, then  $c = c_k^+ = c_k^+(a, b)$  for some  $k$  or  $c = c_j^- = c_j^-(a, b)$  for some  $j$ . The set  $c_j^-$  is omitted if  $b = a + 2$ .*

However, it is difficult to apply this theorem in many cases, since the upper bound of  $b$  is small. Hence, there were a lot of researches to generalize the result. The following lemma generalizes of the Theorem 2.2.

**Lemma 2.3** ([10, Lemma 4.1]). *Let  $\{a, b, c\}$  be a Diophantine triple. Assume that  $a < b \leq 8a$ . Then  $c = c_\nu^\tau$  for some  $\nu$  and  $\tau$ .*

Next, the following theorem gives us the bound of third element  $c$  in the Diophantine triple  $\{a, b, c\}$ .

**Theorem 2.4** ([10, Theorem 1.2]). *Let  $\{a, b, c\}$  be a Diophantine triple with  $a < b$ . Suppose that  $\{a, b, c, d\}$  is a Diophantine quadruple with  $d > d_+$  and that  $\{a, b, c', c\}$  is not a Diophantine quadruple for any  $c'$  with  $0 < c' < d_-$ , where  $d_+$  and  $d_-$  are defined by*

$$d_\pm = a + b + c + 2abc \pm 2rst,$$

*respectively.*

- (1) *If  $b < 2a$ , then  $c < b^6$ .*
- (2) *If  $2a \leq b \leq 8a$ , then  $c < 9.5b^4$ .*
- (3) *If  $b > 8a$ , then  $c < b^5$ .*

If  $c = c_\nu^\tau$ , then we can find the upper bound of  $c$  more specific by the following theorem.

**Theorem 2.5** ([11, Theorem 1.4]). *Suppose that  $\{a, b, c_\nu^\tau, d\}$  is a Diophantine quadruple with  $d > c_{\nu+1}^\tau$  and that  $\{a, b, c', c_\nu^\tau\}$  is not a Diophantine quadruple for any  $c'$  with  $0 < c' < c_{\nu-1}^\tau$ .*

- (1) *If  $b < 2a$ , then  $c \leq c_3^+$ .*
- (2) *If  $2a \leq b \leq 8a$ , then  $c \leq c_2^+$ .*

## 2.2. The Properties of solutions of Pell equation

We have to solve the system

$$ad + 1 = x^2, \quad bd + 1 = y^2, \quad cd + 1 = z^2$$

to extend the Diophantine triple  $\{a, b, c\}$  to the Diophantine quadruple  $\{a, b, c, d\}$ . One can eliminate  $d$  to obtain the following system of Pell equations

- (1)  $ay^2 - bx^2 = a - b,$
- (2)  $az^2 - cx^2 = a - c,$
- (3)  $bz^2 - cy^2 = b - c.$

**Lemma 2.6** ([6, Lemma 1]). *There exist positive integers  $i_0, j_0$  and integers  $z_0^{(i)}, x_0^{(i)}, z_1^{(j)}, y_1^{(j)}, i = 1, \dots, i_0, j = 1, \dots, j_0$ , with the following properties:*

- (1)  *$(z_0^{(i)}, x_0^{(i)})$  and  $(z_1^{(j)}, y_1^{(j)})$  are solutions of (2) and (3), respectively.*
- (2)  *$z_0^{(i)}, x_0^{(i)}, z_1^{(j)}, y_1^{(j)}$  satisfy the following inequalities*

$$0 < x_0^{(i)} \leq \sqrt{\frac{a(c-a)}{2(s-1)}} < \sqrt{\frac{s+1}{2}} < \sqrt[4]{ac},$$

$$0 \leq |z_0^{(i)}| \leq \sqrt{\frac{(s-1)(c-a)}{2a}} < \sqrt{\frac{c\sqrt{c}}{2\sqrt{a}}} < \frac{c}{2},$$

$$0 < y_1^{(j)} \leq \sqrt{\frac{b(c-b)}{2(t-1)}} < \sqrt{\frac{t+1}{2}} < \sqrt[4]{bc},$$

$$0 \leq |z_1^{(j)}| \leq \sqrt{\frac{(t-1)(c-b)}{2b}} < \sqrt{\frac{c\sqrt{c}}{2\sqrt{b}}} < \frac{c}{3}.$$

- (3) *If  $(z, x)$  and  $(z, y)$  are positive integers of (2) and (3), respectively, then there exist  $i \in \{1, \dots, i_0\}, j \in \{1, \dots, j_0\}$  and integers  $m, n \geq 0$  such that*

$$(4) \quad z\sqrt{a} + x\sqrt{c} = (z_0^{(i)}\sqrt{a} + x_0^{(i)}\sqrt{c})(s + \sqrt{ac})^m,$$

$$(5) \quad z\sqrt{b} + y\sqrt{c} = (z_1^{(j)}\sqrt{b} + y_1^{(j)}\sqrt{c})(t + \sqrt{bc})^n.$$

From now on, we omit the superscripts  $(i)$  and  $(j)$ . By (4), we may write  $z = v_m$ , where

$$(6) \quad v_0 = z_0, \quad v_1 = sz_0 + cx_0, \quad v_{m+2} = 2sv_{m+1} - v_m,$$

and by (5), we may write  $z = w_n$ , where

$$(7) \quad w_0 = z_1, \quad w_1 = tz_1 + cy_1, \quad w_{n+2} = 2tw_{n+1} - w_n.$$

**Lemma 2.7** ([8, Lemma 3]). *If  $v_m = w_n$ , then  $n - 1 \leq m \leq 2n + 1$ .*

### 2.3. Congruence relation between solutions of Pell equations

In this section, we give the congruence relations between  $v_m$  and  $w_n$ , and properties of initial terms of (6) and (7).

**Lemma 2.8** ([6, Lemma 4]). *We have the following properties of  $v_m$  and  $w_n$ .*

$$\begin{aligned} v_{2m} &\equiv z_0 + 2c[az_0m^2 + sx_0m] \pmod{8c^2}, \\ v_{2m+1} &\equiv sz_0 + c[2asx_0m(m+1) + x_0(2m+1)] \pmod{4c^2}, \\ w_{2n} &\equiv z_1 + 2c[bz_1n^2 + ty_1n] \pmod{8c^2}, \\ w_{2n+1} &\equiv tz_1 + c[2btz_1n(n+1) + y_1(2n+1)] \pmod{4c^2}. \end{aligned}$$

We have a question such that when does  $v_m = w_n$  have a solution and if there exists a solution of  $v_m = w_n$ , then what are the values possible for the solution. The following lemma gives us the answer.

**Lemma 2.9** ([8, Lemma 8]). *We have the following results.*

- (1) *If the equation  $v_{2m} = w_{2n}$  has a solution, then  $z_0 = z_1$ . Furthermore,  $|z_0| = 1$  or  $|z_0| = cr - st$  or  $|z_0| < \min\{0.869a^{-5/14}c^{9/14}, 0.972b^{-0.3}c^{0.7}\}$ .*
- (2) *If the equation  $v_{2m+1} = w_{2n}$  has a solution, then  $|z_0| = t, |z_1| = cr - st$  and  $z_0z_1 < 0$ .*
- (3) *If the equation  $v_{2m} = w_{2n+1}$  has a solution, then  $|z_0| = cr - st, |z_1| = s$  and  $z_0z_1 < 0$ .*
- (4) *If the equation  $v_{2m+1} = w_{2n+1}$  has a solution, then  $|z_0| = t, |z_1| = s$  and  $z_0z_1 > 0$ .*

Furthermore, the solution of  $v_m = w_n$  is more specific when  $c = c_\nu^\mp \leq c_3^+$  by the following lemma.

**Lemma 2.10** ([11, Lemma 3.1]). *Assume that  $a < b \leq 8a$ .*

(i) *Assume that  $b < 3a$ . In the case of  $c = c_1^-$ , we have  $v_{2m+1} \neq w_{2n}$ ,  $v_{2m} \neq w_{2n+1}$  and  $v_{2m+1} \neq w_{2n+1}$ . Moreover, if  $v_{2m} = w_{2n}$ , then  $z_0 = z_1 = 1$ .*

(ii) *In the case of  $c = c_1^+$ , we have  $v_{2m+1} \neq w_{2n}$ ,  $v_{2m} \neq w_{2n+1}$  and  $v_{2m+1} \neq w_{2n+1}$ . Moreover, if  $v_{2m} = w_{2n}$ , then  $z_0 = z_1$  and  $|z_0| = 1$ .*

(iii) *In the case of  $c = c_2^-$ , we have  $v_{2m+1} \neq w_{2n}$  and  $v_{2m+1} \neq w_{2n+1}$ . Moreover, we have the following:*

- (1) *If  $v_{2m} = w_{2n}$ , then  $z_0 = z_1$  and  $|z_0| = 1$  or  $cr - st$ .*
- (2) *If  $v_{2m} = w_{2n+1}$ , then  $|z_0| = cr - st$  and  $|z_1| = s$  with  $z_0z_1 < 0$ .*

Furthermore, (2) occurs if and only if (1) with  $|z_0| = cr - st$  occurs.

(iv) *In the case of  $c \in \{c_2^+, c_3^-, c_3^+\}$ , we have  $v_{2m+1} \neq w_{2n}$  and  $v_{2m} \neq w_{2n+1}$ . Moreover, we get the following:*

- (1) *If  $v_{2m} = w_{2n}$ , then  $z_0 = z_1$  and  $|z_0| = 1$ .*
- (2) *If  $v_{2m+1} = w_{2n+1}$ , then  $|z_0| = t$  and  $|z_1| = s$  with  $z_0z_1 > 0$ .*

#### 2.4. Some theorems for applying the reduction method

From (4), (5) and sum of their conjugate, respectively, we get

$$v_m = \frac{1}{2\sqrt{a}}[(z_0\sqrt{a} + x_0\sqrt{c})(s + \sqrt{ac})^m + (z_0\sqrt{a} - x_0\sqrt{c})(s - \sqrt{ac})^m],$$

$$w_n = \frac{1}{2\sqrt{b}}[(z_1\sqrt{b} + y_1\sqrt{c})(t + \sqrt{bc})^n + (z_1\sqrt{b} - y_1\sqrt{c})(t - \sqrt{bc})^n].$$

Hence, we transform the equation  $v_m = w_n$  into the following inequality.

**Lemma 2.11** ([6, Lemma 5]). *Assume that  $c > 4b$ . If  $v_m = w_n$  and  $m, n \neq 0$ , then*

$$0 < m \log(s + \sqrt{ac}) - n \log(t + \sqrt{bc}) + \log \frac{\sqrt{b}(x_0\sqrt{c} + z_0\sqrt{a})}{\sqrt{a}(y_1\sqrt{c} + z_1\sqrt{b})} < \frac{8}{3} ac(s + \sqrt{ac})^{-2m}.$$

We use the following theorem and lemma to obtain the upper bound for  $m$ .

**Theorem 2.12** ([2, p. 20]). *For a linear form*

$$\Lambda = \beta_1 \log \alpha_1 + \cdots + \beta_l \log \alpha_l \neq 0$$

*in logarithms of  $l$  algebraic numbers  $\alpha_1, \alpha_2, \dots, \alpha_l$  with rational coefficients  $\beta_1, \beta_2, \dots, \beta_l$ , we have*

$$\log |\Lambda| \geq -18(l+1)!^{l+1} (32d)^{l+2} h'(\alpha_1) \cdots h'(\alpha_l) \log(2ld) \log \beta,$$

*where  $\beta := \max\{|\beta_1|, \dots, |\beta_l|\}$ ,  $d := [\mathbb{Q}(\alpha_1, \dots, \alpha_l) : \mathbb{Q}]$  and*

$$h'(\alpha) = \frac{1}{d} \max\{h(\alpha), |\log \alpha|, 1\}$$

*with the standard logarithmic Weil height  $h(\alpha)$  of  $\alpha$ .*

**Lemma 2.13** ([9, Lemma 5]). *Suppose that  $M$  is a positive integer. Let  $p/q$  be the convergent of the continued fraction expansion of  $\kappa$  such that  $q > 6M$  and let  $\epsilon = \|\mu q\| - M \cdot \|\kappa q\|$ , where  $\|\cdot\|$  denotes the distance from the nearest integer.*

(1) *If  $\epsilon > 0$ , then there is no solution of the inequality*

$$(8) \quad 0 < m\kappa - n + \mu < AB^{-m}$$

*in integers  $m$  and  $n$  with*

$$\frac{\log(Aq/\epsilon)}{\log B} \leq m \leq M.$$

(2) *Let  $r = \lfloor \mu q + \frac{1}{2} \rfloor$ . If  $p - q + r = 0$ , then there is no solution of inequality (8) in integers  $m$  and  $n$  with*

$$\max \left\{ \frac{\log(3Aq)}{\log B}, 1 \right\} < m \leq M.$$

### 3. The generalized bound for the second element

We can easily know that the form of third elements of Diophantine triple  $\{F_{k-2}F_{k+1}, F_{k-1}F_{k+2}, c\}$  by using Lemma 2.3. However, we can't apply the Lemma 2.3 in the case of  $\{F_{k-2}F_{k-1}, F_{k+1}F_{k+2}, c\}$ , since

$$8F_{k-2}F_{k-1} < F_{k+1}F_{k+2} < 21F_{k-2}F_{k-1}.$$

Hence, we need to generalize the Lemma 2.3 to find the form of third elements of Diophantine triple  $\{a, b, c\}$  when  $b > 8a$ .

**Lemma 3.1.** *Let  $\{a, b, c\}$  be a Diophantine triple and  $a < b \leq 24a$ . Suppose that  $\{1, 3, a, b\}$  is not a Diophantine quadruple. Then  $c = c_v^\tau$  for some  $n$  and  $\tau$ .*

*Proof.* The proof is same as the proof of Lemma 4.1 in [10], except the case of  $c' < b$ . If  $c' = 0$ , then  $s' = rs - at = 1$ , and  $c = c_1^-$  or  $c_1^+$ . Let  $r' = s'r - at'$  and  $b' = ((r')^2 - 1)/a$ . Then,  $b' = a + b + c' + 2abc' - 2rs't'$ , and thus, [15, Lemma 4] and  $b \leq 24a$  together imply that

$$b' < \frac{b}{4ac'} \leq \frac{24a}{4ac'} = \frac{6}{c'}.$$

First, we consider the case of  $c' = 1$ . Since  $b'$  and  $c'$  satisfy the condition  $b'c' + 1 = b' + 1 = \square$ , we have  $b' = 0$ . For the other cases, we have the following results.

$$b' = \begin{cases} 0 & \text{if } c' \geq 6, \\ 0, 1 & \text{if } c' = 3, 4, 5, \\ 0, 1, 2 & \text{if } c' = 2. \end{cases}$$

We need to check only the case  $(b', c') = (1, 3)$  except  $b' = 0$  using the condition  $b'c' + 1 = \square$ . However, this means the set  $\{1, 3, a, b\}$  satisfies Diophantine quadruple, which is a contradiction by assumption. In all cases, we obtain  $b' = 0$  which yields  $c' = a + b - 2r = c_1^-$  and  $c = c_2^-$ . Hence, we proved the lemma.  $\square$

Assume that  $a < b \leq 24a$  and  $\{1, 3, a, b\}$  is not a Diophantine quadruple. Then, Lemma 3.1 implies that the positive solutions of the Diophantine equation

$$ay^2 - bx^2 = a - b$$

are given by

$$y\sqrt{a} + x\sqrt{b} = (\pm\sqrt{a} + \sqrt{b})(r + \sqrt{ab})^l$$

with some non-negative integer  $l$ . Thus, we may write  $x = p_l$ ,  $y = V_l$ , where

$$(9) \quad p_0 = 1, \quad p_1 = r \pm a, \quad p_{l+2} = 2rp_{l+1} - p_l,$$

$$(10) \quad V_0 = \pm 1, \quad V_1 = b \pm r, \quad V_{l+2} = 2rV_{l+1} - V_l.$$

From equation (4) and (5), we may also write  $x = q_m$ ,  $y = W_n$ , where

$$(11) \quad q_0 = x_0, \quad q_1 = sx_0 + az_0, \quad q_{m+2} = 2sq_{m+1} - q_m,$$

$$(12) \quad W_0 = y_1, \quad W_1 = ty_1 + bz_1, \quad W_{n+2} = 2tW_{n+1} - W_n.$$



The result of next lemma is similar to the Lemma 2.10, but the bound of  $b$  is further extended.

**Lemma 3.2.** *Assume that  $a < b \leq 24a$  and  $\{1, 3, a, b\}$  is not a Diophantine quadruple.*

(i) *In the case of  $c_1^+$ , we have  $v_{2m+1} \neq w_{2n}$ ,  $v_{2m} \neq w_{2n+1}$  and  $v_{2m+1} \neq w_{2n+1}$ . Moreover, if  $v_{2m} = w_{2n}$ , then  $z_0 = z_1$  and  $|z_0| = 1$ .*

(ii) *In the case of  $c = c_2^-$ , we have  $v_{2m+1} \neq w_{2n}$  and  $v_{2m+1} \neq w_{2n+1}$ . Moreover, we have the following:*

(1) *If  $v_{2m} = w_{2n}$ , then  $z_0 = z_1$  and  $|z_0| = 1$  or  $cr - st$ .*

(2) *If  $v_{2m} = w_{2n+1}$ , then  $|z_0| = cr - st$  and  $|z_1| = s$  with  $z_0 z_1 < 0$ .*

*Furthermore, (2) occurs if and only if (1) with  $|z_0| = cr - st$  occurs.*

(iii) *In the case of  $c \in \{c_2^+, c_3^-, c_3^+\}$ , then we have  $v_{2m+1} \neq w_{2n}$  and  $v_{2m} \neq w_{2n+1}$ . Moreover, we get the following:*

• *If  $v_{2m} = w_{2n}$ , then  $z_0 = z_1$  and  $|z_0| = 1$ .*

• *If  $v_{2m+1} = w_{2n+1}$ , then  $|z_0| = t$  and  $|z_1| = s$  with  $z_0 z_1 > 0$ .*

*Proof.* The proof proceeds along the same lines as that of [11, Lemma 3.1].

(i) From (10) and (12), we have  $y_1^2 \equiv 1 \pmod{b}$  either  $n$  is even or odd in  $W_n$ . Since  $a < b - 2$  and  $r < b - 1$ , so  $c = c_1^+ = a + b + 2r \leq 4b - 7$ . Hence

$$0 < y_1^2 \leq \frac{t+1}{2} \leq \frac{\sqrt{b(4b-7)+1}}{2} < b.$$

This means  $y_1^2 = 1$ , that is,  $y_1 = 1$  and  $z_1 = \pm 1$ . Hence, the case  $v_{2m} = w_{2n}$  only occur and  $z_0 = z_1$  from Lemma 2.9.

(ii) Suppose that  $|z_0| = t$ . Then we have  $c > 4ab^2$  by upper bound of  $y$ . However, this is a contradiction, since

$$c = c_2^- = 4ab \left\{ b - \left( 2r - a - \frac{1}{a} \right) - \frac{r-a}{ab} \right\} < 4ab^2.$$

Therefore, we have  $|z_0| \neq t$  and this means  $v_{2m+1} \neq w_{2n}$  and  $v_{2m+1} \neq w_{2n+1}$  by Lemma 2.9. Also, the last case of  $|z_0|$  in  $v_{2m} = w_{2n}$  occurs when the Diophantine quadruple  $\{a, b, c, d\}$  is only irregular. Hence, it suffices to show that any Diophantine quadruple  $\{a, b, c, d_0\}$  with  $d_0 < c$  is regular when the cases of  $c \leq c_3^+$ . Assume that the set  $\{a, b, c, d_0\}$  with  $d_0 < d_-$  is a Diophantine quadruple. First, since  $r = \sqrt{ab+1} \leq \sqrt{24a^2+1} < 5a$  and together with Lemma 2.1, we have following inequality

$$c \leq c_3^+ < a^{2.5} b^{3.5} \left( \frac{176}{b} + \frac{16}{b^{0.5}} + \frac{184}{b^{2.5}} + \frac{24}{b^{1.5}} + \frac{39}{b^{3.5}} + \frac{9}{b^{2.5}} \right) < a^{2.5} b^{3.5}.$$

However, we have  $c > 16a^{2.5} b^{3.5}$  by Lemma 15 of [14]. Hence, Diophantine quadruple  $\{a, b, c, d_0\}$  with  $d_0 < c$  can not be irregular. It is easy to see that  $v_{2m}$  and  $w_{2n+1}$  in (2) are equal to  $v_{2m}$  and  $w_{2n}$  in (1) with  $|z_0| = cr - st$ .

(iii) Since we show that if  $c \leq c_3^+$ , then any Diophantine quadruple  $\{a, b, c, d_0\}$  with  $d_0 < c$  is regular in (ii), it suffices to show that  $\max\{|z_0|, |z_1|\} < cr - st$  when  $c \geq c_2^+$ . Since  $c \geq c_2^+ > 4ab^2 \geq 32ab$ , we have

$$cr - st = \frac{c^2 - (a+b)c - 1}{cr + st} > \frac{(1 - a/c - b/c - 1/c^2)c}{2\sqrt{ab}\sqrt{1 + 1/(ab)}} > \frac{c}{2.1\sqrt{ab}}.$$

If  $\max\{|z_0|, |z_1|\} \geq cr - st$ , then the bounds of  $|z_0|$  and  $|z_1|$  show that

$$c < \left(\frac{2.1}{\sqrt{2}}\right)^4 ab^2 < 5ab^2.$$

However, we have the following inequalities

$$c \geq c_2^+ > 4ab(a+b+2r) > 4ab^2 \left(\frac{1}{24} + 1 + \frac{1}{\sqrt{6}}\right) > 5.79ab^2,$$

since  $b \leq 24a$  implies that  $r = \sqrt{ab+1} > b/(2\sqrt{6})$ . Hence, this is a contradiction.  $\square$

#### 4. The extendibility of $\{F_{k-2}F_{k+1}, F_{k-1}F_{k+2}\}$

In this section, we prove the extendibility of the Diophantine pair  $\{F_{k-2}F_{k+1}, F_{k-1}F_{k+2}\}$ . From Theorem 2.5, we only need to check the cases of  $c \leq c_2^+$ .

**Lemma 4.1.** *Suppose that  $m, n \geq 2$ .*

(1) *If  $v_{2m} = w_{2n}$ , then*

$$m \geq \begin{cases} (\sqrt{2F_k+1} - 1)/2, & \text{if } c_1^+ \text{ or } c_2^- \text{ with } |z_0| = cr - st, \\ \sqrt[4]{F_k}/2, & \text{if } c_2^- \text{ with } |z_0| = 1 \text{ or } c_2^+. \end{cases}$$

(2) *If  $v_{2m} = w_{2n+1}$  with  $c_2^-$  or  $v_{2m+1} = w_{2n+1}$  with  $c_2^+$ , then*

$$m \geq \frac{\sqrt{2F_k+1} - 1}{2}.$$

*Proof.* Since  $c_i^\pm$  is divisible by  $F_k$  for all  $i = 1, 2$  and

- In the case of  $c_1^+$ ,  
 $s_1^+ \equiv F_{k-2}F_{k+1} = a \pmod{F_k}$  and  
 $t_1^+ \equiv -F_{k-2}F_{k+1} = -a \pmod{F_k}$ .
- In the case of  $c_2^-$ ,  
 $s_2^- = 2F_{k-1}^2F_k^2 - 1 \equiv -1 \pmod{F_k}$  and  
 $t_2^- = 2F_k^2(F_{k-1}F_k \pm 1) + 1 \equiv 1 \pmod{F_k}$ .
- In the case of  $c_2^+$ ,  
 $s_2^+ = 2F_{k-2}F_{k+1}^3 + 1 \equiv -1 \pmod{F_k}$  and  
 $t_2^+ = 2F_k^2F_{k+1}^2 - 1 \equiv -1 \pmod{F_k}$ .

Hence, we have the following lower bounds from Lemma 2.8.

(1) The case of  $v_{2m} = w_{2n}$ :

We easily find the relation between  $v_{2m}$  and  $w_{2n}$  such that

$$\pm am^2 + sm \equiv \pm bn^2 + tn \pmod{c}.$$

- First, we consider the case of  $c_1^+$ .

Since  $F_{k-1}F_{k+2} \equiv -F_{k-2}F_{k+1} \pmod{F_k}$  and  $\gcd(F_{k-2}F_{k+1}, F_k) = 1$ , so we have

$$\pm(m^2 + n^2 \pm m \pm n) \equiv 0 \pmod{F_k}.$$

This means  $2(m^2 + m) \geq m^2 + n^2 \pm m \pm n \geq m^2 + n^2 - m - n > 0$ .  
Hence  $2(m^2 + m) \geq F_k$ .

- In the case of  $c_2^-$  with  $|z_0| = cr - st$ , we get the equation

$$\pm am^2 + am \equiv \pm bn^2 + bn \pmod{F_k},$$

since  $st \equiv -1 \pmod{F_k}$  and  $r \equiv 0 \pmod{F_k}$ . Hence, similar to above, we have the result.

- Let us consider the other case of  $c_2^-$ , that is,  $c_2^-$  with  $|z_0| = 1$ . We have the equation

$$\pm a(m^2 + n^2) \equiv m + n \pmod{F_k},$$

and squaring both sides, then  $a^2(m^2 + n^2)^2 \equiv (m + n)^2 \pmod{F_k}$ .  
Since  $a^2 \equiv 1 \pmod{F_k}$ , we have

$$(m^2 + n^2)^2 - (m + n)^2 \geq 0.$$

This means  $4m^4 \geq F_k$ , so we have the desired result.

- Lastly, in the case of  $c_2^+$  with  $|z_0| = 1$ , we have

$$\pm a(m^2 + n^2) \equiv m - n \pmod{F_k}.$$

Similar to above, squaring both sides and we have

$$(m^2 + n^2) - (m - n)^2 \equiv 0 \pmod{F_k}.$$

Hence,  $4m^4 \geq F_k$ , and the result is deduced.

(2) The cases of  $v_{2m} = w_{2n+1}$  with  $c_2^-$  and  $v_{2m+1} = w_{2n+1}$  with  $c_2^+$ :  
From Lemma 2.8, we have

$$\pm a(-st)m^2 - astm \equiv \mp 2bstn(n+1) \pmod{F_k}$$

and

$$\pm 2atsm(m+1) \equiv \pm 2bstn(n+1) \pmod{F_k},$$

respectively. So, they become

$$a(m^2 + n^2 + n \pm m) \equiv 0 \pmod{F_k}$$

and

$$\pm a(m^2 + n^2 + m + n) \equiv 0 \pmod{F_k},$$

respectively, and we have the desired result.  $\square$

Let us find the logarithmic inequality for  $c = c_1^+$ .

**Lemma 4.2.** *If  $v_{2m} = w_{2n}$  with  $c_1^+$  and  $m, n \neq 0$ , then*

$$\begin{aligned} 0 &< 2m \log(s + \sqrt{ac}) - 2n \log(t + \sqrt{bc}) + \log \frac{\sqrt{b}(x_0\sqrt{c} + z_0\sqrt{a})}{\sqrt{a}(y_1\sqrt{c} + z_1\sqrt{b})} \\ &< 3.018(s + \sqrt{ac})^{-4m}. \end{aligned}$$

*Proof.* Put

$$P = \frac{1}{\sqrt{a}}(x_0\sqrt{c} + z_0\sqrt{a})(s + \sqrt{ac})^m, \quad Q = \frac{1}{\sqrt{b}}(y_1\sqrt{c} + z_1\sqrt{b})(t + \sqrt{bc})^n.$$

Then

$$P^{-1} = \frac{\sqrt{a}(x_0\sqrt{c} - z_0\sqrt{a})}{c - a}(s - \sqrt{ac})^m, \quad Q^{-1} = \frac{\sqrt{b}(y_1\sqrt{c} - z_1\sqrt{b})}{c - b}(t - \sqrt{bc})^n.$$

Therefore, the relation  $v_m = w_n$  becomes

$$P - \frac{c-a}{a}P^{-1} = Q - \frac{c-b}{b}Q^{-1}.$$

Since  $P > 0$ ,  $Q > 0$  and

$$P - Q > \frac{c-a}{a}(Q - P)P^{-1}Q^{-1},$$

it follows that  $P > Q$ . Furthermore, we have

$$\frac{P-Q}{P} < \frac{c-a}{a}P^{-2} < \frac{1}{a(c-a)} \leq \frac{1}{170},$$

since the case of  $\{3, 5\}$  is proved by Filipin, Fujita and Togbé [11]. Hence,

$$\begin{aligned} 0 &< \log \frac{P}{Q} = -\log\left(1 - \frac{P-Q}{P}\right) < \frac{171}{170} \left(\frac{c-a}{a}\right)P^{-2} \\ &< \frac{171}{170} \frac{c-a}{(\sqrt{c}-\sqrt{a})^2} (s + \sqrt{ac})^{-2m}. \end{aligned}$$

Since  $\frac{\sqrt{c}+\sqrt{a}}{\sqrt{c}-\sqrt{a}} < 3$ , we obtain the result.  $\square$

#### 4.1. The Theorem of Baker and Wüstholz

Now we apply theorem of Baker and Wüstholz.

(i) First, we consider the equation  $v_{2m} = w_{2n}$ . We may assume that  $k > 2$ . Using Lemma 2.11 and Lemma 4.2, and apply Theorem 2.12, then we have  $l = 3$ ,  $d = 4$ ,  $\beta = 2m$ ,

$$\alpha_1 = s + \sqrt{ac}, \quad \alpha_2 = t + \sqrt{bc}, \quad \alpha_3 = \frac{(\sqrt{c} \pm \sqrt{a})\sqrt{b}}{(\sqrt{c} \pm \sqrt{b})\sqrt{a}}.$$

Let  $\alpha_3'$  and  $\alpha_3''$  be the conjugates of  $\alpha_3$  whose absolute values are greater than one. Then

$$h'(\alpha_1) = \frac{1}{2} \log(\alpha_1) < \frac{1}{2} \log(2s), \quad h'(\alpha_2) = \frac{1}{2} \log(\alpha_2) < \frac{1}{2} \log(2t),$$

$$\begin{aligned} h'(\alpha_3) &\leq \frac{1}{4} \{ \log(a^2(c-b)^2) + \log(\alpha_3 \alpha_3' \alpha_3'') \} \\ &= \frac{1}{4} \{ \log(b\sqrt{ab}(\sqrt{c} + \sqrt{a})(\sqrt{c} + \sqrt{b})(c-a)) \} < \log(1.42c), \end{aligned}$$

and

$$\log |\Lambda| \geq -18 \cdot 4! 3^4 (32 \cdot 4)^5 \frac{1}{2} \log(2s) \frac{1}{2} \log(2t) \log(1.42c) \cdot \log(24) \cdot \log(2m).$$

Since

$$\log\left(\frac{8}{3} ac(s + \sqrt{ac})^{-4m}\right) < (-2m + 1) \log(4ac)$$

and

$$\log(3.018(s + \sqrt{ac})^{-4m}) < (-2m + 1) \log(4ac),$$

we have

$$(13) \quad \frac{2m-1}{\log(2m)} < 9.556 \cdot 10^{14} \log(2c) \log(1.42c).$$

Let  $x = F_k$ .

- If  $c = c_1^+$ , then

$$\sqrt{2x+1} - 2 < 4.778 \cdot 10^{14} \log(3.17x)^2 \cdot \log(2.67x)^2 \cdot \log(2x+1).$$

Hence,  $x < 1.64 \cdot 10^{42}$  and  $c_1^+ < 1.35 \cdot 10^{85}$ .

- If  $c = c_2^-$  with  $|z_0| = cr - st$ , then

$$\sqrt{2x+1} - 2 < 4.778 \cdot 10^{14} \log(2x)^6 \cdot \log(1.89x)^6 \cdot \log(2x+1).$$

Hence,  $x < 1.72 \cdot 10^{44}$  and  $c_2^- < 6.22 \cdot 10^{266}$ .

- If  $c = c_2^-$  with  $|z_0| = \pm 1$ , then

$$\sqrt[4]{x} - 1 < 2.389 \cdot 10^{14} \log(2x)^6 \cdot \log(1.89x)^6 \cdot \log x.$$

Hence,  $x < 4.38 \cdot 10^{91}$  and  $c_2^- < 1.68 \cdot 10^{551}$ .

- If  $c = c_2^+$ , then

$$\sqrt[4]{x} - 1 < 2.389 \cdot 10^{14} \log(2.55x)^6 \cdot \log(2.41x)^6 \cdot \log x.$$

Hence,  $x < 4.42 \cdot 10^{91}$  and  $c_2^+ < 8.83 \cdot 10^{551}$ .

Since  $F_k = (\alpha^k - \bar{\alpha}^k)/\sqrt{5}$ , where  $\alpha = (1 + \sqrt{5})/2 > 1.618$ , we have the following inequality from Fibonacci numbers

$$(14) \quad (1.618)^k < (\alpha)^k = \bar{\alpha}^k + \sqrt{5} \cdot F_k.$$

We can find the upper bounds of  $k$  and  $m$  by using inequalities (13) and (14), respectively.

- If the case of  $c_1^+$ , then  $k \leq 203$  and  $m \leq 9.04 \cdot 10^{20}$ .
- If the case of  $c_2^-$ , then  $k \leq 440$  and  $m \leq 4.07 \cdot 10^{22}$ .
- If the case of  $c_2^+$ , then  $k \leq 440$  and  $m \leq 4.08 \cdot 10^{22}$ .

(ii) Let  $v_{2m} = w_{2n+1}$  with  $n \neq 0$ . We have  $l = 3$ ,  $d = 4$ ,  $\beta = 2m + 1$ ,

$$\alpha_1 = s + \sqrt{ac}, \quad \alpha_2 = t + \sqrt{bc}, \quad \alpha_4 = \frac{((sr - ta)\sqrt{c} \pm (cr - st)\sqrt{a})\sqrt{b}}{(r\sqrt{c} \mp s\sqrt{b})\sqrt{a}}.$$

Let  $\alpha'_4$  and  $\alpha''_4$  be the conjugates of  $\alpha_4$  whose absolute values are greater than one. Then

$$\begin{aligned} h'(\alpha_1) &= \frac{1}{2} \log(\alpha_1) < \frac{1}{2} \log(2s), \quad h'(\alpha_2) = \frac{1}{2} \log(\alpha_2) < \frac{1}{2} \log(2t), \\ h'(\alpha_4) &\leq \frac{1}{4} \{\log(a^2(c-b)^2) + \log(\alpha_4 \alpha'_4 \alpha''_4)\} \\ &= \frac{1}{4} \{\log(b\sqrt{ab}((sr - ta)\sqrt{c} + (cr - st)\sqrt{a})(r\sqrt{c} + s\sqrt{b})(c-a))\} < \log(1.42\sqrt{rc}), \end{aligned}$$

since  $(sr - ta)\sqrt{c} > (cr - st)\sqrt{a}$  and  $sr - ta < r$ . Hence, we have

$$(15) \quad \frac{2m-1}{\log(2m+1)} < 9.556 \cdot 10^{14} \log(2c) \log(1.42\sqrt{rc}).$$

If  $c = c_2^-$ , then

$$\sqrt{x+1} - 2 < 4.778 \cdot 10^{14} \log(1.97x)^6 \cdot \log(1.7x)^7 \cdot \log(x+1),$$

where  $x = F_k$ . Hence, we have  $x < 4.9 \cdot 10^{44}$  and  $c_2^- < 3.98 \cdot 10^{269}$ . Again, by using the inequality from (14) and (15), we get the bound of  $k \leq 215$  and  $m \leq 1.11 \cdot 10^{22}$ .

(iii) Let  $v_{2m+1} = w_{2n+1}$  with  $n \neq 0$ . We have  $l = 3$ ,  $d = 4$ ,  $\beta = 2m + 1$ ,

$$\alpha_1 = s + \sqrt{ac}, \quad \alpha_2 = t + \sqrt{bc}, \quad \alpha_5 = \frac{(r\sqrt{c} \pm t\sqrt{a})\sqrt{b}}{(r\sqrt{c} \pm s\sqrt{b})\sqrt{a}}.$$

Let  $\alpha'_5$  and  $\alpha''_5$  be the conjugates of  $\alpha_5$  whose absolute values are greater than one. Then

$$\begin{aligned} h'(\alpha_1) &= \frac{1}{2} \log(\alpha_1) < \frac{1}{2} \log(2s), \quad h'(\alpha_2) = \frac{1}{2} \log(\alpha_2) < \frac{1}{2} \log(2t), \\ h'(\alpha_5) &\leq \frac{1}{4} \{\log(a^2(c-b)^2) + \log(\alpha_5 \alpha'_5 \alpha''_5)\} \\ &= \frac{1}{4} \{\log(b\sqrt{ab}(r\sqrt{c} + t\sqrt{a})(r\sqrt{c} + s\sqrt{b})(c-a))\} < \log(1.42\sqrt{rc}). \end{aligned}$$

Hence, we have

$$(16) \quad \frac{2m}{\log(2m+1)} < 9.556 \cdot 10^{14} \log(2c) \log(1.42\sqrt{rc}).$$

If  $c = c_2^+$ , then

$$\sqrt{x+1} - 1 < 4.778 \cdot 10^{14} \log(2.53x)^6 \cdot \log(2.11x)^7 \cdot \log(x+1),$$

where  $x = F_{2k+2}$ . Hence, we get  $x < 4.95 \cdot 10^{44}$  and  $c_2^+ < 1.91 \cdot 10^{270}$ . This means  $k \leq 215$  and  $m \leq 1.12 \cdot 10^{22}$ .

#### 4.2. The reduction method

Now dividing logarithmic inequalities from Lemma 2.11 and Lemma 4.2 by  $\log \alpha_2$ , respectively leads us to the inequalities

$$\begin{aligned} 0 &< m_1\kappa - n_1 + \mu_1 < A_1B^{m_1}, \\ 0 &< m_1\kappa - n_1 + \mu_1 < A_2B^{m_1}, \\ 0 &< m_1\kappa - n_2 + \mu_2 < A_1B^{m_1}, \\ 0 &< m_2\kappa - n_2 + \mu_3 < A_1B^{m_2}, \end{aligned}$$

where  $m_1 := 2m$ ,  $m_2 := 2m + 1$ ,  $n_1 := 2n$ ,  $n_2 := 2n + 1$  and

$$\begin{aligned} \kappa &= \frac{\log \alpha_1}{\log \alpha_2}, \quad \mu_1 = \frac{\log \alpha_3}{\log \alpha_2}, \quad \mu_2 = \frac{\log \alpha_4}{\log \alpha_2}, \quad \mu_3 = \frac{\log \alpha_5}{\log \alpha_2}, \\ A_1 &= \frac{(8/3)ac}{\log \alpha_2}, \quad A_2 = \frac{3.018}{\log \alpha_2}, \quad B = \alpha_1^2. \end{aligned}$$

We apply Lemma 2.13 to the logarithmic inequalities with  $M_1 := 2m \leq 8.16 \cdot 10^{22}$  and  $M_2 := 2m + 1 \leq 2.24 \cdot 10^{22}$ . We have to examine  $2 \cdot 203 + 2 \cdot 440 + 2 \cdot 440 + 2 \cdot 215 + 2 \cdot 215 = 3026$  cases. The program was developed in **PARI/GP** running with 400 digits. For the computations, if the first convergent such that  $q > 6M_i$  with  $i = 1, 2$  does not satisfy the condition  $\epsilon > 0$ , then we use the next convergent until we find the one that satisfies the conditions. Then we have the results as the following Table 1.

TABLE 1. Results from **PARI/GP** running

Case of $c$	Initial values	Use the next convergent
$c_1^+$	$z_0 = z_1 = 1$	0 case
	$z_0 = z_1 = -1$	99 cases
$c_2^-$	$z_0 = z_1 = 1$	330 cases
	$z_0 = z_1 = -1$	352 cases ( $k = 89, \dots, 440$ )
$c_2^-$	$z_0 = cr - st, z_1 = -s$	0 case
	$z_0 = st - cr, z_1 = s$	0 case
$c_2^+$	$z_0 = z_1 = 1$	353 cases ( $k = 88, \dots, 440$ )
	$z_0 = z_1 = -1$	353 cases ( $k = 88, \dots, 440$ )
$c_2^+$	$z_0 = t, z_1 = s$	0 case
	$z_0 = -t, z_1 = -s$	172 cases ( $k = 44, \dots, 215$ )

We have the upper bounds 13, 12 and 8 of  $m$  in the case of  $c_1^+$ ,  $c_2^-$  with  $|z_0| = 1$ , and  $c_2^+$  with  $|z_0| = 1$ , respectively. If we take  $M = 2m$  and continue the program again, then we have  $m \leq 1$ . Other cases, that is,  $c_2^-$  with  $|z_0| = cr - st, |z_1| = s$ , and  $c_2^+$  with  $|z_0| = t, |z_1| = s$  have the upper bounds 7 and 8 of

$m$ , respectively. Hence, we take  $M = 2m$  or  $2m + 1$ , and continue the program again, we also get  $m \leq 1$ . Therefore, we get the following theorem.

**Theorem 4.3.** *Let  $k \geq 3$  be an integer and the set  $\{F_{k-2}F_{k+1}, F_{k-1}F_{k+2}, c, d\}$  be a Diophantine quadruple with  $c < d$ , then  $d = d_+$ .*

**Corollary 4.4.** *The set  $\{F_{k-2}F_{k+1}, F_{k-1}F_{k+2}, c\}$  can be extended only to regular.*

### 5. The extendibility of $\{F_{k-2}F_{k-1}, F_{k+1}F_{k+2}\}$

In this section, we show that the set  $\{F_{k-2}F_{k-1}, F_{k+1}F_{k+2}, c\}$  can be extended to only regular, where  $c > \max\{F_{k-2}F_{k-1}, F_{k+1}F_{k+2}\}$ . We can easily show that the Fibonacci numbers  $F_{k-2}F_{k-1}$  and  $F_{k+1}F_{k+2}$  do not satisfy the Diophantine quadruple with  $\{1, 3\}$ , by considering the period of Fibonacci numbers in modulo 4. Hence, we can find the form of third element in the Diophantine triple  $\{F_{k-2}F_{k-1}, F_{k+1}F_{k+2}, c\}$  by Lemma 3.1.

**Lemma 5.1.** *Let  $\{F_{k-2}F_{k-1}, F_{k+1}F_{k+2}, c\}$  be a Diophantine triple. Then  $c = c_\nu^\tau$  for some  $\nu$  and  $\tau$ .*

The small values of  $c_\nu^\tau$  are listed below.

$$\begin{aligned} c_1^+ &= 4F_k F_{k+1}, \\ c_2^- &= 4F_k^2(4F_k^3 F_{k-1} + 1), \\ c_2^+ &= 4F_k^2(4F_k^3 F_{k+1} - 1), \\ c_3^- &= F_k[64F_{k-2}^2 F_{k-1}^3 F_{k+1}^2 F_{k+2}^2 + 16F_{k-2} F_{k-1} F_k F_{k+1} F_{k+2} \\ &\quad + 96F_{k-2} F_{k-1}^2 F_{k+1} F_{k+2} + 12F_k + 36F_{k-1}], \\ c_3^+ &= F_k[64F_{k-2}^2 F_{k-1}^2 F_{k+1}^3 F_{k+2}^2 + 96F_{k-2} F_{k-1} F_k F_{k+1}^2 F_{k+2} \\ &\quad - 16F_{k-2} F_{k-1}^2 F_{k+1} F_{k+2} + 24F_k + 36F_{k-1}]. \end{aligned}$$

By the Theorem 2.4 and Lemma 3.1, we have the upper bound of third elements  $c$  in the Diophantine triple  $\{a, b, c\}$  with  $a < b \leq 21a$ .

**Lemma 5.2.** *Let  $\{a, b\}$  be a Diophantine pair with  $a < b \leq 21a$ . Suppose that  $\{a, b, c, d\}$  is a Diophantine quadruple with  $d > c_{n+1}^\tau$  and that  $\{a, b, c', c\}$  is not a Diophantine quadruple for any  $c'$  with  $0 < c' < c_{n-1}^\tau$ . Then  $c \leq c_3^+$ .*

*Proof.* Since  $c_4^- = 64a^3 b^3(a + b - 2r) + 64a^2 b^2(2a + 2b - 3r) + 80ab(a + b - r) + 8(2a + 2b - r)$ , it suffices to show that  $64a^3 b^3(a + b - 2r) > b^5$ , by Theorem 2.4. We have

$$\frac{c_4^-}{b^5} > \frac{64a^3(a + b - 2r)}{b^2} > \frac{64a^3(a + b - 2r)}{(21a)^2} > 1.16 > 1.$$

Hence, this completes the proof of lemma.  $\square$

**Lemma 5.3.** *Suppose that  $m, n \geq 2$ .*



(1) If  $v_{2m} = w_{2n}$ , then

$$m \geq \begin{cases} (\sqrt[4]{F_k}/2, & \text{if } c_2^- \text{ with } |z_0| = 1 \text{ or } c_2^+, \\ \sqrt{2F_k + 1} - 1)/2, & \text{if the other cases.} \end{cases}$$

(2) If  $v_{2m} = w_{2n+1}$  with  $c_2^-$  or  $v_{2m+1} = w_{2n+1}$  with  $c_2^+$ ,  $c_3^-$  and  $c_3^+$ , then

$$m \geq \frac{\sqrt{2F_k + 1} - 1}{2}.$$

*Proof.* We have the following relation between  $c_\nu^r$  and  $F_k$ .

- In the case of  $c_1^+$ :  $s_1^+ \equiv a \pmod{F_k}$ ,  $t_1^+ \equiv b \equiv -a \pmod{F_k}$ .
- In the case of  $c_2^-$ :  $s_2^- \equiv -1 \pmod{F_k}$ ,  $t_2^- \equiv 1 \pmod{F_k}$ .
- In the case of  $c_2^+$ :  $s_2^+ \equiv -1 \pmod{F_k}$ ,  $t_2^+ \equiv 1 \pmod{F_k}$ .
- In the case of  $c_3^-$ :  $s_3^- \equiv a \pmod{F_k}$ ,  $t_3^- \equiv -b \equiv a \pmod{F_k}$ .
- In the case of  $c_3^+$ :  $s_3^+ \equiv -a \pmod{F_k}$ ,  $t_3^+ \equiv -b \equiv a \pmod{F_k}$ .

We can also easily find the relation between  $v_m$  and  $w_n$ .

(1) The case of  $v_{2m} = w_{2n}$ :

$$\pm am^2 + sm \equiv \pm bn^2 + tn \pmod{F_k}.$$

(2) The case of  $v_{2m} = w_{2n+1}$ :

$$\pm a(-st)m^2 - astm \equiv \mp 2bstn(n+1) \pmod{F_k}.$$

(3) The case of  $v_{2m+1} = w_{2n+1}$ :

$$\pm 2atsm(m+1) \equiv \pm 2bstn(n+1) \pmod{F_k}.$$

Since  $F_{k+1}F_{k+2} \equiv -F_{k-2}F_{k-1} \pmod{F_k}$  and  $\gcd(F_{k-2}F_{k-1}, F_k) = 1$ , we can prove the lemma by the same procedure as in the proof of Lemma 4.1.  $\square$

**Lemma 5.4.** If  $v_{2m} = w_{2n}$  with  $c_1^+$  and  $m, n \neq 0$ , then

$$\begin{aligned} 0 &< 2m \log(s + \sqrt{ac}) - 2n \log(t + \sqrt{bc}) + \log \frac{\sqrt{b}(x_0\sqrt{c} + z_0\sqrt{a})}{\sqrt{a}(y_1\sqrt{c} + z_1\sqrt{b})} \\ &< 2.09(s + \sqrt{ac})^{-4m}. \end{aligned}$$

*Proof.* The proof follows along the same line as that of Lemma 4.2. In this case, we have

$$\frac{P-Q}{P} < \frac{c-a}{a}P^{-2} < \frac{1}{a(c-a)} \leq \frac{1}{23}.$$

Hence,

$$0 < \log \frac{P}{Q} = -\log\left(1 - \frac{P-Q}{P}\right) < \frac{24}{23} \left(\frac{c-a}{a}\right)P^{-2} < \frac{24}{23} \frac{c-a}{(\sqrt{c}-\sqrt{a})^2} (s+\sqrt{ac})^{-2m}.$$

Since  $\frac{\sqrt{c}+\sqrt{a}}{\sqrt{c}-\sqrt{a}} < 2$ , we obtain the result.  $\square$

### 5.1. The Theorem of Baker and Wüstholz

By the theorem of Baker and Wüstholz again, we have the equations (13), (15) and (16) for  $v_{2m} = w_{2n}$ ,  $v_{2m} = w_{2n+1}$  and  $v_{2m+1} = w_{2n+1}$ , respectively, since  $2.09(s + \sqrt{ac})^{-4m} < (-2m + 1) \log(4ac)$ .

Let  $x = F_k$ . First, let us consider the cases of the equation  $v_{2m} = w_{2n}$ . We have the following inequalities for each  $c_i^\pm$ ,  $i = 1, 2, 3$ .

- If  $c = c_1^+$ , then

$$\sqrt{2x+1} - 2 < 4.778 \cdot 10^{14} \log(4x)^2 \cdot \log(3.38x)^2 \cdot \log(2x+1).$$

Hence,  $x < 1.65 \cdot 10^{42}$  and  $c_1^+ < 2.18 \cdot 10^{85}$ .

- If  $c = c_2^-$ , then

$$\sqrt[4]{x} - 1 < 2.389 \cdot 10^{14} \log(1.85x)^6 \cdot \log(1.75x)^6 \cdot \log x.$$

Hence,  $x < 4.36 \cdot 10^{91}$  and  $c_2^- < 1.1 \cdot 10^{551}$ .

- If  $c = c_2^+$ , then

$$\sqrt[4]{x} - 1 < 2.389 \cdot 10^{14} \log(2.04x)^6 \cdot \log(1.93x)^6 \cdot \log x.$$

Hence,  $x < 4.38 \cdot 10^{91}$  and  $c_2^+ < 2.26 \cdot 10^{551}$ .

- If  $c = c_3^-$ , then

$$\sqrt{2x+1} - 2 < 4.778 \cdot 10^{14} \log(2.39x)^{10} \cdot \log(2.31x)^{10} \cdot \log(2x+1).$$

Hence,  $x < 1.52 \cdot 10^{45}$  and  $c_3^- < 2 \cdot 10^{455}$ .

- If  $c = c_3^+$ , then

$$\sqrt{2x+1} - 2 < 4.778 \cdot 10^{14} \log(2.56x)^{10} \cdot \log(2.47x)^{10} \cdot \log(2x+1).$$

Hence,  $x < 1.52 \cdot 10^{45}$  and  $c_3^+ < 3.84 \cdot 10^{455}$ .

From (14), we get the upper bound of  $k$  for each  $c$ . Also from (13) and the upper bound of  $c$ , we get the upper bound of  $m$ .

- If  $c = c_1^+$ , then  $k \leq 203$  and  $m \leq 9.08 \cdot 10^{20}$ .
- If  $c = c_2^-$ , then  $k \leq 440$  and  $m \leq 4.07 \cdot 10^{22}$ .
- If  $c = c_2^+$ , then  $k \leq 440$  and  $m \leq 4.07 \cdot 10^{22}$ .
- If  $c = c_3^-$ , then  $k \leq 217$  and  $m \leq 2.76 \cdot 10^{22}$ .
- If  $c = c_3^+$ , then  $k \leq 217$  and  $m \leq 2.76 \cdot 10^{22}$ .

Next, we consider the case of  $v_{2m} = w_{2n+1}$  with  $c_2^\pm$  with  $|z_0| = cr - st$  and  $|z_1| = s$ . Then we have the inequality from (15).

$$\sqrt{2x+1} - 2 < 4.778 \cdot 10^{14} \log(1.85x)^6 \cdot \log(1.62x)^7 \cdot \log(x+1).$$

Hence,  $x < 2.34 \cdot 10^{44}$  and  $c_2^- < 3.29 \cdot 10^{267}$ . This means  $k \leq 213$  and  $m \leq 1.09 \cdot 10^{22}$ .

Lastly, consider the case of  $v_{2m+1} = w_{2n+1}$ . There are three cases  $c_2^+$ ,  $c_3^-$  and  $c_3^+$ . We have the following inequalities from (16).

- In the case of  $c = c_2^+$ , then  
 $\sqrt{x+1} - 1 < 4.778 \cdot 10^{14} \log(2.04x)^6 \cdot \log(1.76x)^7 \cdot \log(x+1)$ .  
Hence,  $x < 4.91 \cdot 10^{44}$  and  $c_2^+ < 5.05 \cdot 10^{269}$ .
- In the case of  $c = c_3^-$ , then  
 $\sqrt{x+1} - 1 < 4.778 \cdot 10^{14} \log(2.39x)^{10} \cdot \log(2.14x)^{11} \cdot \log(x+1)$ .  
Hence,  $x < 3.81 \cdot 10^{45}$  and  $c_3^- < 1.95 \cdot 10^{459}$ .
- In the case of  $c = c_3^+$ , then  
 $\sqrt{x+1} - 1 < 4.778 \cdot 10^{14} \log(2.56x)^{10} \cdot \log(2.28x)^{11} \cdot \log(x+1)$ .  
Hence,  $x < 3.82 \cdot 10^{45}$  and  $c_3^+ < 3.86 \cdot 10^{459}$ .

Also, using the inequality (14) and upper bounds of  $c$ , we have the following upper bound of  $k$  and  $m$  for each cases.

- In the case of  $c = c_2^+$ , then  $k \leq 215$  and  $m \leq 1.11 \cdot 10^{22}$ .
- In the case of  $c = c_3^-$ , then  $k \leq 219$  and  $m \leq 3.09 \cdot 10^{22}$ .
- In the case of  $c = c_3^+$ , then  $k \leq 219$  and  $m \leq 3.09 \cdot 10^{22}$ .

## 5.2. The reduction method

We again apply Lemma 2.13 about diving logarithmic inequalities from Lemma 2.11 and Lemma 5.4 by  $\log \alpha_2$ , that is,

$$\begin{aligned} 0 &< m_1 \kappa - n_1 + \mu_1 < A_1 B^{m_1}, \\ 0 &< m_1 \kappa - n_1 + \mu_1 < A_2 B^{m_1}, \\ 0 &< m_1 \kappa - n_2 + \mu_2 < A_1 B^{m_1}, \\ 0 &< m_2 \kappa - n_2 + \mu_3 < A_1 B^{m_2}, \end{aligned}$$

where  $m_1 := 2m$ ,  $m_2 := 2m + 1$ ,  $n_1 := 2n$ ,  $n_2 := 2n + 1$  and

$$\begin{aligned} \kappa &= \frac{\log \alpha_1}{\log \alpha_2}, \quad \mu_1 = \frac{\log \alpha_3}{\log \alpha_2}, \quad \mu_2 = \frac{\log \alpha_4}{\log \alpha_2}, \quad \mu_3 = \frac{\log \alpha_5}{\log \alpha_2}, \\ A_1 &= \frac{(8/3)ac}{\log \alpha_2}, \quad A_2 = \frac{2.09}{\log \alpha_2}, \quad B = \alpha_1^2. \end{aligned}$$

Let  $M_1 := 2m \leq 5.52 \cdot 10^{22}$  and  $M_2 := 2m + 1 \leq 6.18 \cdot 10^{22}$ . We have to examine  $2 \cdot 203 + 2 \cdot 440 + 2 \cdot 440 + 2 \cdot 217 + 2 \cdot 217 + 2 \cdot 213 + 2 \cdot 215 + 2 \cdot 219 + 2 \cdot 219 = 4766$  cases. Similar to Section 4.2.2, we have the result as the following Table 2 by using the program **PARI/GP** running with 400 digit. We have the following upper bounds of  $m$  in Table 2 for each cases. Use these upper bounds and continue the program again, then we have  $m \leq 1$ . Hence, we get the following theorem.

**Theorem 5.5.** *Let  $k \geq 3$  be the integer and the set  $\{F_{k-2}F_{k-1}, F_{k+1}F_{k+2}, c, d\}$  be a Diophantine quadruple with  $c < d$ , then  $d = d_+$ .*

**Corollary 5.6.** *The set  $\{F_{k-2}F_{k-1}, F_{k+1}F_{k+2}, c\}$  can be extended only to regular.*

TABLE 2. Results from PARI/GP running

Case of $c$	Initial values	Use the next convergent	Upper bound of $m$
$c_1^+$	$z_0 = z_1 = 1$	0 case	15
	$z_0 = z_1 = -1$	0 case	14
$c_2^-$	$z_0 = z_1 = 1$	353 cases ( $k = 88, \dots, 440$ )	10
	$z_0 = z_1 = -1$	353 cases ( $k = 88, \dots, 440$ )	10
$c_2^-$	$z_0 = cr - st, z_1 = -s$	169 cases ( $k = 45, \dots, 213$ )	10
	$z_0 = st - cr, z_1 = s$	0 case	10
$c_2^+$	$z_0 = z_1 = 1$	354 cases ( $k = 86, 88, \dots, 440$ )	9
	$z_0 = z_1 = -1$	0 case	8
$c_2^+$	$z_0 = t, z_1 = s$	200 cases ( $k = 16, \dots, 215$ )	8
	$z_0 = -t, z_1 = -s$	171 cases ( $k = 45, \dots, 215$ )	8
$c_3^-$	$z_0 = z_1 = 1$	0 case	6
	$z_0 = z_1 = -1$	0 case	6
$c_3^-$	$z_0 = t, z_1 = s$	204 cases ( $k = 12, \dots, 215$ )	6
	$z_0 = -t, z_1 = -s$	0 case	7
$c_3^+$	$z_0 = z_1 = 1$	174 cases ( $k = 44, \dots, 217$ )	6
	$z_0 = z_1 = -1$	0 case	6
$c_3^+$	$z_0 = t, z_1 = s$	0 case	6
	$z_0 = -t, z_1 = -s$	197 cases ( $k = 23, \dots, 219$ )	6

### References

- [1] A. Baker and H. Davenport, *The equations  $3x^2 - 2 = y^2$  and  $8x^2 - 7 = z^2$* , Quart. J. Math. Oxford Ser. (2) **20** (1969), 129–137.
- [2] A. Baker and G. Wüstholz, *Logarithmic forms and group varieties*, J. Reine Angew. Math. **442** (1993), 19–62.
- [3] Y. Bugeaud, A. Dujella, and M. Mignotte, *On the family of Diophantine triples  $\{k - 1, k + 1, 16k^3 - 4k\}$* , Glasg. Math. J. **49** (2007), no. 2, 333–344.
- [4] A. Dujella, *The problem of the extension of a parametric family of Diophantine triples*, Publ. Math. Debrecen **51**, (1997), no. 3-4, 311–322.
- [5] ———, *A parametric family of elliptic curves*, Acta Arith. **94**, (2000), no. 1, 87–101.
- [6] ———, *An absolute bound for the size of Diophantine  $m$ -tuple*, J. Number theory **89** (2001), no. 1, 126–150.
- [7] ———, *Diophantine  $m$ -tuples and elliptic curves*, 21st Journées Arithmétiques (Rome, 2001), J. Théor. Nombres Bordeaux **13**, (2001), no. 1, 111–124.
- [8] ———, *There are only finitely many Diophantine quintuples*, J. Reine Angew. Math. **566** (2004), 183–214.
- [9] A. Dujella and A. Pethö, *Generalization of a theorem of Baker and Davenport*, Quart. J. Math. Oxford Ser. (2) **49** (1998), no. 195, 291–306.
- [10] A. Filipin, Y. Fujita, and A. Togbé, *The extendibility of Diophantine pairs I: The general case*, Glas. Mat. Ser. III **49**(69) (2014), no. 1, 25–36.
- [11] ———, *The extendibility of Diophantine pairs II: Examples*, J. Number Theory **145** (2014), 604–631.

- [12] Y. Fujita, *The extensibility of Diophantine pair  $\{k - 1, k + 1\}$* , J. Number Theory **128** (2008), no. 2, 323–353.
- [13] ———, *The Hoggatt-Bergum conjecture on  $D(-1)$ -triples  $\{F_{2k+1}, F_{2k+3}, F_{2k+5}\}$  and integer points on the attached elliptic curves*, Rocky Mountain J. Math. **39** (2009), no. 6, 1907–1932.
- [14] ———, *Any Diophantine quintuple contains a regular Diophantine quadruple*, J. Number Theory **129** (2009), no. 7, 1678–1697.
- [15] B. W. Jones, *A second variation on a problem of Diophantus and Davenport*, Fibonacci Quart. **16** (1978), no. 2, 155–165.

JUNE BOK LEE  
DEPARTMENT OF MATHEMATICAL SCIENCES  
YONSEI UNIVERSITY  
SEOUL **Please supply Zip Code**, KOREA  
*E-mail address:* leejb@yonsei.ac.kr

JINSEO PARK  
DEPARTMENT OF MATHEMATICAL SCIENCES  
YONSEI UNIVERSITY  
SEOUL **Please supply Zip Code**, KOREA  
*E-mail address:* dbjin2000@yonsei.ac.kr