

# GLOBAL NONEXISTENCE FOR THE WAVE EQUATION WITH BOUNDARY VARIABLE EXPONENT NONLINEARITIES

TAE GAB HA AND SUN-HYE PARK

**ABSTRACT.** This paper deals with a nonlinear wave equation with boundary damping and source terms of variable exponent nonlinearities. This work is devoted to prove a global nonexistence of solutions for a nonlinear wave equation with nonnegative initial energy as well as negative initial energy.

## 1. Introduction

In this paper, we consider the following the wave equation:

$$(1) \quad \begin{cases} u_{tt} - \mu(t)\Delta u + h(u) = 0 & \text{in } \Omega \times (0, T), \\ u = 0 & \text{on } \Gamma_0 \times (0, T), \\ \mu(t)\frac{\partial u}{\partial \nu} + |u_t|^{m(x)-2}u_t = |u|^{p(x)-2}u & \text{on } \Gamma_1 \times (0, T), \\ u(x, 0) = u_0(x), \quad u'(x, 0) = u_1(x) & \text{in } \Omega, \end{cases}$$

where,  $\Omega$  is a bounded open domain of  $\mathbb{R}^n (n \geq 1)$  with a smooth boundary  $\Gamma = \Gamma_0 \cup \Gamma_1$ . Here,  $\Gamma_0$  and  $\Gamma_1$  are closed and disjoint with  $meas(\Gamma_0) > 0$ . Let  $\nu$  be the outward normal to  $\Gamma$  and  $T > 0$ , a real number, and  $m(x)$ ,  $p(x)$  be given functions.

This type of model arises in electro-rheological fluids or fluids with temperature dependent viscosity, viscoelasticity, filtration processes through a porous media and image processing (cf. [1, 20]).

The problem of proving the nonexistence or blow-up of solutions for the wave equation has been widely studied (see [6, 8–13, 17, 18, 21–23]). Recently, many papers have treated problems with variable exponents. For the variable

---

Received May 30, 2021; Revised September 2, 2021; Accepted September 27, 2021.

2010 *Mathematics Subject Classification.* Primary 35L05, 35A01, 35B44.

*Key words and phrases.* Global nonexistence, boundary variable exponent nonlinearity.

T. G. Ha was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (NRF-2019R1I1A3A01051714). S.-H. Park was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (2020R1I1A3066250).

exponent problems, the main tool is based on the Lebesgue and Sobolev spaces with variable exponents, which was introduced in [3, 4] and has been widely used in the literature, see [2, 7, 14–16] and the list of references therein. For example, in [16], the authors proved the local existence of a unique weak solution for the nonlinear damped wave equation and the finite time blow-up of solutions for negative initial energies. Recently, in [7], the authors studied the global existence of solution for (1) using the stable-set method and proved the exponential or polynomial energy decay rate. However, the above mentioned references was only considered interior variable exponent nonlinearities.

On the other hand, there are very few results for the boundary variable-exponent-nonlinearity problems. In [19], the author proved the existence and asymptotic stability for the semilinear wave equation with boundary variable exponent nonlinearities. However, the blow-up was not considered.

Motivated by previous works, the goal of this paper is to prove a finite time blow-up for the solution for (1) under suitable condition on the initial data and the positive initial energy. As far as we know, there is no blow-up result concerning the boundary variable-exponent nonlinearities.

This paper is organized as follows: In Section 2, we recall the notation, hypotheses and some necessary preliminaries and introduce our main result. In Section 3, we prove the blow-up of solutions for (1) with nonnegative initial energy as well as negative initial energy.

## 2. Preliminaries

We begin this section by introducing some hypotheses and our main result. Throughout this paper,  $\|\cdot\|_p$  and  $\|\cdot\|_{p,\Gamma_1}$  denote the  $L^p(\Omega)$  norm and  $L^p(\Gamma_1)$  norm, respectively.

### (H<sub>1</sub>) Hypotheses on $\Omega$ .

Let  $\Omega \subset \mathbb{R}^n$  be a bounded open domain,  $n \geq 1$ , with a smooth boundary  $\Gamma = \Gamma_0 \cup \Gamma_1$ . Here  $\Gamma_0$  and  $\Gamma_1$  are closed and disjoint with  $meas(\Gamma_0) > 0$ , satisfying the following conditions:

$$(2) \quad \begin{aligned} w(x) \cdot \nu(x) &\geq \sigma > 0 \quad \text{on } \Gamma_1, \quad w(x) \cdot \nu(x) \leq 0 \quad \text{on } \Gamma_0, \\ w(x) &= x - x^0 \quad (x^0 \in \mathbb{R}^n) \quad \text{and} \quad R = \max_{x \in \bar{\Omega}} |w(x)|, \end{aligned}$$

where  $\nu$  represents the unit outward normal vector to  $\Gamma$ . We assume that

$$(3) \quad \mu(0) \frac{\partial u_0}{\partial \nu} + |u_1|^{m(x)-2} u_1 = |u_0|^{p(x)-2} u_0 \quad \text{on } \Gamma_1.$$

### (H<sub>2</sub>) Hypotheses on $m(x)$ , $p(x)$ .

Let  $m(x)$  and  $p(x)$  be given measurable functions on  $\bar{\Omega}$  satisfying the following conditions:

$$(4) \quad \begin{cases} 2 \leq q^- \leq q(x) \leq q^+ < \frac{2(n-1)}{n-2} & \text{if } n \geq 3, \\ q^- > 2 & \text{if } n = 1, 2, \end{cases}$$

where

$$q^- = \operatorname{ess\,inf}_{x \in \Omega} q(x), \quad \text{and} \quad q^+ = \operatorname{ess\,sup}_{x \in \Omega} q(x).$$

Furthermore,  $m(x)$  and  $p(x)$  satisfy the log-Hölder continuity condition as follows

$$(5) \quad |q(x) - q(y)| \leq -\frac{A}{\log|x-y|} \quad \text{for all } x, y \in \Omega,$$

with  $|x - y| < \delta$ ,  $A > 0$ ,  $0 < \delta < 1$ .

**(H<sub>3</sub>) Hypotheses on  $\mu$ ,  $h$ .**

Let  $\mu \in W^{1,\infty}(0, \infty) \cap W^{1,1}(0, \infty)$  satisfy the following conditions:

$$(6) \quad \mu(t) \geq \mu_0 > 0 \quad \text{and} \quad \mu'(t) < 0 \quad \text{a.e. in } [0, \infty),$$

where  $\mu_0$  is a positive constant. Moreover, we assume that

$$(7) \quad h : \mathbb{R} \rightarrow \mathbb{R} \text{ is a Lipschitz function, and } 2H(s) \geq h(s)s \geq 0 \text{ for all } s \in \mathbb{R},$$

where  $H(s) = \int_0^s h(\tau) d\tau$ .

In order to treat the variable-exponent nonlinearities  $m(x)$  and  $p(x)$ , we need some preliminary facts about the Lebesgue and Sobolev spaces with variable exponents (see [3, 4]). For the reader's convenience, we will repeat some of them here.

Let  $q : \Omega \rightarrow [1, \infty]$  be a measurable function. We define the Lebesgue space with a variable exponent  $q(\cdot)$  by

$$L^{q(\cdot)}(\Omega) := \{u \mid u : \Omega \rightarrow \mathbb{R} \text{ measurable and } \int_{\Omega} |\lambda u(x)|^{q(x)} dx < \infty \text{ for some } \lambda > 0\},$$

equipped with the following Luxembourg-type norm

$$\|u\|_{q(\cdot), \Omega} = \|u\|_{q(\cdot)} := \inf\{\lambda > 0 \mid \int_{\Omega} \left| \frac{u(x)}{\lambda} \right|^{q(x)} dx \leq 1\}.$$

$L^{q(\cdot)}(\Omega)$  is a Banach space. Next, we define the Sobolev space  $W^{1,q(\cdot)}(\Omega)$  as follows:

$$W^{1,q(\cdot)}(\Omega) := \{u \in L^{q(\cdot)}(\Omega) \text{ such that } \nabla u \text{ exists and } |\nabla u| \in L^{q(\cdot)}(\Omega)\}.$$

This is a Banach space with respect to the norm  $\|u\|_{W^{1,q(\cdot)}(\Omega)} = \|u\|_{q(\cdot)} + \|\nabla u\|_{q(\cdot)}$ . Furthermore, we set  $W_0^{1,q(\cdot)}(\Omega)$  to be the closure of  $C_0^\infty(\Omega)$  in the space  $W^{1,q(\cdot)}(\Omega)$ .

**Lemma 2.1** (Poincaré's Inequality [3, 4]). *Let  $\Omega$  be a bounded domain of  $\mathbb{R}^n$  and  $q(\cdot)$  satisfies (5). Then*

$$\|u\|_{q(\cdot)} \leq C_1 \|\nabla u\|_{q(\cdot)} \quad \text{for all } u \in W_0^{1,q(\cdot)}(\Omega),$$

where  $C_1$  is a positive constant which depends on  $q^\pm$  and  $\Omega$ . In particular,  $\|\nabla u\|_{q(\cdot)}$  defines an equivalent norm on  $W_0^{1,q(\cdot)}(\Omega)$ .

**Lemma 2.2** ([3,4]). *Let  $\Omega$  be a bounded domain of  $\mathbb{R}^n$  with a smooth boundary  $\Gamma$ . Assume that  $r : \Omega \rightarrow (1, \infty)$  is a measurable function such that*

$$1 < r^- \leq r(x) \leq r^+ < +\infty \quad \text{for a.e. } x \in \Omega.$$

*If  $q(x), r(x) \in C(\bar{\Omega})$  and  $q(x) < r^*(x)$  in  $\bar{\Omega}$  with*

$$r^*(x) = \begin{cases} \frac{nr(x)}{n-r(x)} & \text{if } r^+ < n, \\ \infty & \text{if } r^+ \geq n. \end{cases}$$

*Then the embedding  $W^{1,r(\cdot)}(\Omega) \hookrightarrow L^{q(\cdot)}(\Omega)$  is continuous and compact.*

**Lemma 2.3** ([5]). *Let  $\Omega$  be a bounded domain in  $\mathbb{R}^n$ ,  $q \in C^{0,1}(\bar{\Omega})$ ,  $1 < q^- \leq q(x) \leq q^+ < n$ . Then for any  $r \in C(\Gamma)$  with  $1 \leq r(x) \leq \frac{(n-1)q(x)}{n-q(x)}$ , there is a continuous trace  $W^{1,q(\cdot)}(\Omega) \hookrightarrow L^{r(\cdot)}(\Gamma)$ , and when  $1 \leq r(x) < \frac{(n-1)q(x)}{n-q(x)}$ , the trace is compact, in particular, the continuous trace  $W^{1,q(\cdot)}(\Omega) \hookrightarrow L^{q(\cdot)}(\Gamma)$  is compact.*

From Lemmas 2.2 and 2.3, we have the embedding  $H_0^1(\Omega) \hookrightarrow L^{q(\cdot)}(\Gamma_1)$ , where

$$H_0^1(\Omega) = \{u \in H^1(\Omega) : u = 0 \text{ on } \Gamma_1\}$$

and

$$\begin{cases} 2 \leq q^- \leq q(x) \leq q^+ < \frac{2(n-1)}{n-2} & \text{if } n \geq 3, \\ q^- > 2 & \text{if } n = 1, 2, \end{cases}$$

which satisfies the inequalities

$$(8) \quad \|u\|_{q(\cdot)} \leq C_2 \|\nabla u\|_2 \quad \text{and} \quad \|u\|_{q(\cdot), \Gamma_1} \leq C_3 \|\nabla u\|_2 \quad \text{for all } u \in H_0^1(\Omega),$$

where  $C_2$  and  $C_3$  are for some positive constants.

**Lemma 2.4** (Hölder's inequality [3,4]). *Let  $q, r, s \geq 1$  be measurable functions defined on  $\Omega$  such that*

$$\frac{1}{s(x)} = \frac{1}{q(x)} + \frac{1}{r(x)} \quad \text{for a.e. } x \in \Omega.$$

*If  $f \in L^{q(\cdot)}(\Omega)$  and  $g \in L^{r(\cdot)}(\Omega)$ , then  $fg \in L^{s(\cdot)}(\Omega)$ ,*

$$\int_{\Omega} |fg|^s(x) dx \leq \int_{\Omega} |f|^q(x) dx + \int_{\Omega} |g|^r(x) dx$$

*and*

$$\|fg\|_{s(\cdot)} \leq 2 \|f\|_{q(\cdot)} \|g\|_{r(\cdot)}.$$

**Lemma 2.5** ([3,4]). *If  $q : \bar{\Omega} \rightarrow [1, \infty)$  is a continuous function satisfying  $2 \leq q_1 \leq q(x) \leq q_2 < q^*$ , where*

$$\begin{cases} q^* = \frac{2n}{n-2} & \text{for } n \geq 3, \\ q^* = \infty & \text{for } n = 1, 2. \end{cases}$$

Then the embedding  $H_0^1(\Omega) \hookrightarrow L^{q(\cdot)}(\Omega)$  is continuous and compact, and we have

$$(9) \quad \min \left\{ \|u\|_{q(\cdot), \Omega}^{q_1}, \|u\|_{q(\cdot), \Omega}^{q_2} \right\} \leq \int_{\bar{\Omega}} |u|^{q(x)} dx \leq \max \left\{ \|u\|_{q(\cdot), \Omega}^{q_1}, \|u\|_{q(\cdot), \Omega}^{q_2} \right\}.$$

The following theorem is the local existence of solution of problem (1), which can be established employing the Faedo-Galerkin method as in the work of [16, 19].

**Theorem 2.6.** *Let the initial data  $\{u_0, u_1\}$  belong to  $H_0^1(\Omega) \times L^2(\Omega)$  and the hypotheses  $(H_1)$ – $(H_3)$  hold. Additionally  $p(x)$  satisfies*

$$(10) \quad 2 \leq p^- \leq p(x) \leq p^+ < \frac{2n-3}{n-2} \quad \text{if } n \geq 3.$$

Then problem (1) has a unique weak solution such that

$$u \in L^\infty((0, T); H_0^1(\Omega)), \quad L^\infty((0, T); L^2(\Omega)) \cap L^{m(\cdot)}(\Gamma_1 \times (0, T)).$$

In order to formulate another result, it is convenient to introduce the energy associated with problem (1):

$$(11) \quad E(t) = \frac{1}{2} \|u_t\|_2^2 + \frac{1}{2} \mu(t) \|\nabla u\|_2^2 + \int_{\Omega} H(u) dx - \int_{\Gamma_1} \frac{|u|^{p(x)}}{p(x)} d\Gamma,$$

where  $H(s) = \int_0^s h(\tau) d\tau$ . Then by (6),

$$E'(t) = \frac{1}{2} \mu'(t) \|\nabla u\|_2^2 - \int_{\Gamma_1} |u_t|^{m(x)} d\Gamma \leq 0,$$

which implies that  $E(t)$  is a nonincreasing function.

**Theorem 2.7.** *Suppose that the hypotheses  $(H_1)$ – $(H_3)$  hold. Moreover, we assume that  $m^+ < p^-$  and,*

$$(12) \quad E(0) < d \quad \text{and} \quad \eta_1 < \|\nabla u_0\|_2 \leq C_4^{-1},$$

where  $d = \mu_0 \left( \frac{1}{2} - \frac{1}{p^-} \right) \eta_1^2$ ,  $\eta_1 = (\mu_0 C_4^{-p^-})^{\frac{1}{p^- - 2}}$  and  $C_4 = \max\{1, C_3\}$ . Then the solution of problem (1) cannot exist for all time.

### 3. Blow-up

This section is devoted to prove Theorem 2.7. By similar arguments as in [16] and using (8), we get the following lemma.

**Lemma 3.1.** *Suppose that the assumption (6) holds and  $u$  is a solution of (1). Then we have*

$$(13) \quad \left( \int_{\Gamma_1} |u|^{p(x)} d\Gamma \right)^{\frac{s}{p^-}} \leq C_5 \left( \|\nabla u\|_2^2 + \int_{\Gamma_1} |u|^{p(x)} d\Gamma \right),$$

$$(14) \quad \int_{\Gamma_1} |u|^{p(x)} d\Gamma \geq C_6 \|u\|_{p^-}^{p^-},$$

$$(15) \quad \int_{\Gamma_1} |u|^{m(x)} d\Gamma \leq C_7 \left( \left( \int_{\Gamma_1} |u|^{p(x)} d\Gamma \right)^{\frac{m^-}{p^-}} + \left( \int_{\Gamma_1} |u|^{p(x)} d\Gamma \right)^{\frac{m^+}{p^+}} \right),$$

where  $2 \leq s \leq p^-$  and  $C_5, C_6, C_7 > 1$  are positive constants depending only on  $\Omega$ .

The next lemma plays an essential role for the proof of the blow-up result.

**Lemma 3.2.** *Let the assumption in Theorem 2.7 be satisfied. Then there exists a positive constant  $\eta_*$  such that*

$$(16) \quad \|\nabla u(t)\|_2 \geq \eta_* \quad \text{for all } 0 < t < T_{\max},$$

where  $T_{\max}$  is the maximal time of existence of the solution of (1).

*Proof. Case 1 :*  $0 \leq E(0) < d$ .

By using (11), (6), (9) and (8), we get that

$$(17) \quad \begin{aligned} E(t) &\geq \frac{1}{2} \mu(t) \|\nabla u\|_2^2 - \frac{1}{p^-} \int_{\Gamma_1} |u|^{p(x)} d\Gamma \\ &\geq \frac{1}{2} \mu_0 \|\nabla u\|_2^2 - \frac{1}{p^-} \max \left\{ \|u\|_{p(\cdot), \Gamma_1}^{p^-}, \|u\|_{p(\cdot), \Gamma_1}^{p^+} \right\} \\ &\geq \frac{1}{2} \mu_0 \|\nabla u\|_2^2 - \frac{1}{p^-} \max \left\{ C_3^{p^-} \|\nabla u\|_2^{p^-}, C_3^{p^+} \|\nabla u\|_2^{p^+} \right\} \\ &:= f(\|\nabla u(t)\|_2) \end{aligned}$$

for any  $t \in [0, T_{\max})$ .

We note that  $f(\eta) = g(\eta)$  for  $0 \leq \eta \leq C_4^{-1}$ , where  $g(\eta) = \frac{1}{2} \mu_0 \eta^2 - \frac{C_4^{p^-}}{p^-} \eta^{p^-}$ . It is easy to verify that  $g$  is strictly increasing on  $(0, \eta_1)$  and strictly decreasing on  $(\eta_1, \infty)$ , where  $\eta_1 = (\mu_0 C_4^{-p^-})^{\frac{1}{p^- - 2}}$  is the maximum point of  $g(\eta)$ , and  $g(\eta_1) = d$ . Hence we have  $g(\eta) \rightarrow -\infty$  as  $\eta \rightarrow \infty$ . Since  $E(0) < d = g(\eta_1)$ , there exists  $\eta_2 > \eta_1$  such that  $g(\eta_2) = E(0)$ . Therefore we obtain from (17),

$$g(\eta_2) = E(0) \geq f(\|\nabla u_0\|_2) = g(\|\nabla u_0\|_2),$$

which implies that  $\eta_2 \leq \|\nabla u_0\|_2$ . From (12), we also have

$$(18) \quad \eta_2 \leq C_4^{-1}.$$

Now we prove that

$$(19) \quad \|\nabla u(t)\|_2 \geq \eta_2 \quad \text{for all } 0 < t < T_{\max}$$

by using the contradiction method. Suppose that (19) does not hold. Then there exists  $t^* \in (0, T_{\max})$  such that

$$(20) \quad \|\nabla u(t^*)\|_2 < \eta_2.$$

If  $\|\nabla u(t^*)\|_2 > \eta_1$ , then we obtain from (17), (18) and (20),

$$E(t^*) \geq f(\|\nabla u(t^*)\|_2) = g(\|\nabla u(t^*)\|_2) > g(\eta_2) = E(0),$$

which is a contradiction with respect to the monotonicity of the energy.

If  $\|\nabla u(t^*)\|_2 \leq \eta_1$ , then since  $\eta_1 < \eta_2$ , there exists  $\eta_3$  which verifies

$$\|\nabla u(t^*)\|_2 \leq \eta_1 < \eta_3 < \eta_2 \leq \|\nabla u_0\|_2.$$

From the continuity of the function  $\|\nabla u(\cdot)\|_2$ , there exists  $\bar{t} \in (0, t^*)$  verifying  $\|\nabla u(\bar{t})\|_2 = \eta_3$ . Therefore from (17) and (18) we deduce

$$E(\bar{t}) \geq f(\|\nabla u(\bar{t})\|_2) = g(\|\nabla u(\bar{t})\|_2) > g(\eta_2) = E(0),$$

which is also contradiction.

**Case 2 :**  $E(0) < 0$ .

There is  $\eta_4 > \eta_1$  such that  $g(\eta_4) = E(0)$ , consequently, by (17) we have

$$g(\eta_4) = E(0) \geq f(\|\nabla u_0\|_2) = g(\|\nabla u_0\|_2).$$

From the fact  $g(\eta)$  is decreasing for  $\eta_1 < \eta$ , we get

$$\|\nabla u_0\|_2 \geq \eta_4.$$

By the same argument as in Case 1, we obtain

$$\|\nabla u(t)\|_2 \geq \eta_4 \quad \text{for all } 0 < t < T_{\max}.$$

Let  $\eta_* = \max\{\eta_2, \eta_4\}$ . Then the proof of Lemma 3.2 is completed.  $\square$

Now we will prove the blow-up result in finite time. We set

$$(21) \quad G(t) = E_1 - E(t),$$

where  $E_1$  is a constant lying in  $(E(0), d)$ . Then

$$(22) \quad G'(t) = -E'(t) \geq \int_{\Gamma_1} |u_t|^{m(x)} d\Gamma \geq 0,$$

which implies that  $G(t)$  is a nondecreasing function, consequently, from Lemma 3.2 and the definition of  $d$ , and using the fact that  $\eta_1 < \eta_*$  and  $\mu_0 > 0$ ,

$$(23) \quad 0 < G(0) \leq G(t) < d - \frac{1}{2}\mu_0\eta_*^2 + \frac{1}{p^-} \int_{\Gamma_1} |u|^{p(x)} d\Gamma \leq \frac{1}{p^-} \int_{\Gamma_1} |u|^{p(x)} d\Gamma.$$

We define

$$(24) \quad L(t) = G^{1-\delta}(t) + \epsilon N(t), \quad N(t) = \int_{\Omega} u_t u dx,$$

where  $\epsilon > 0$  will be chosen later and

$$(25) \quad 0 < \delta \leq \min\left\{\frac{p^- - 2}{2p^-}, \frac{p^- - m^+}{p^-(m^+ - 1)}\right\}.$$

Then we have

$$(26) \quad L'(t) = (1 - \delta)G^{-\delta}(t)G'(t) + \epsilon N'(t).$$

We are now going to analyze the last term on the right hand side of (26). From the definition of  $E(t)$ , Lemma 3.2 and (7), we have

$$\begin{aligned}
 N'(t) &= \|u_t\|_2^2 - \mu(t)\|\nabla u\|_2^2 - \int_{\Omega} h(u)u dx + \int_{\Gamma_1} |u|^{p(x)} d\Gamma \\
 &\quad - \int_{\Gamma_1} |u_t|^{m(x)-2} u_t u d\Gamma + \theta E(t) - \theta E(t) \\
 (27) \quad &\geq \left(1 + \frac{\theta}{2}\right) \|u_t\|_2^2 + \underbrace{\mu_0 \left(\frac{\theta}{2} - 1\right) \|\nabla u\|_2^2 - \theta E_1}_{:= I_1} + \theta G(t) \\
 &\quad + \left(1 - \frac{\theta}{p^-}\right) \int_{\Gamma_1} |u|^{p(x)} d\Gamma - \underbrace{\int_{\Gamma_1} |u_t|^{m(x)-2} u_t u d\Gamma}_{:= I_2},
 \end{aligned}$$

provided that  $\theta = p^- - \epsilon$  with  $0 < \epsilon < p^- - 2$ .

**Estimate for  $I_1$ .**

From Lemma 3.2 and the definition of  $\theta$ , we obtain

$$\begin{aligned}
 \mu_0 \left(\frac{\theta}{2} - 1\right) \|\nabla u\|_2^2 - \theta E_1 &> \mu_0 \left(\frac{p^- - \epsilon}{2} - 1\right) \eta_1^2 - (p^- - \epsilon) E_1 \\
 &= \left(E_1 - \frac{\mu_0 \eta_1^2}{2}\right) \epsilon + \mu_0 \left(\frac{p^-}{2} - 1\right) \eta_1^2 - p^- E_1 := F(\epsilon).
 \end{aligned}$$

We note that

$$E_1 - \frac{\mu_0 \eta_1^2}{2} < d - \frac{\mu_0 \eta_1^2}{2} = -\frac{1}{p^-} \mu_0 \eta_1^2 < 0$$

and

$$\mu_0 \left(\frac{p^-}{2} - 1\right) \eta_1^2 - p^- E_1 > \mu_0 \left(\frac{p^-}{2} - 1\right) \eta_1^2 - p^- d = 0.$$

Thus,  $F(\epsilon)$  represent a decreasing line connecting vertical and horizontal axes points  $v_\epsilon := \mu_0 \left(\frac{p^-}{2} - 1\right) \eta_1^2 - p^- E_1$  and  $h_\epsilon := v_\epsilon \left(\frac{\mu_0 \eta_1^2}{2} - E_1\right)^{-1}$ , respectively. Hence, we get that

$$(28) \quad \mu_0 \left(\frac{\theta}{2} - 1\right) \|\nabla u\|_2^2 - \theta E_1 > F(\epsilon) > 0 \quad \text{for } 0 < \epsilon < h_\epsilon.$$


**Estimate for  $I_2$ .**

By multiplying by  $1 = \xi \xi^{-1}$  for  $\xi > 0$ , and by using Lemma 2.4 with  $s = 1$ ,  $q(x) = \frac{m(x)}{m(x)-1}$  and  $r(x) = m(x)$ , it holds that

$$\left| \int_{\Gamma_1} |u_t|^{m(x)-2} u_t u d\Gamma \right| \leq \int_{\Gamma_1} \xi^{m(x)} |u|^{m(x)} d\Gamma + \int_{\Gamma_1} \xi^{-\frac{m(x)}{m(x)-1}} |u_t|^{m(x)} d\Gamma.$$

We take  $\xi^{-\frac{m(x)}{m(x)-1}} = kG^{-\delta}(t)$ , for a large constant  $k$  to be chosen later. The choice of  $\xi$  is allowed since  $G(t) > 0$  for every  $t$  as (23) holds true. Hence the




 FIGURE 1. The figure of  $F(\epsilon)$ 

above inequality takes the form:

$$(29) \quad \int_{\Gamma_1} |u_t|^{m(x)-1} |u| d\Gamma \leq k G^{-\delta}(t) G'(t) + k^{1-m^-} G^{\delta(m^+-1)}(t) \int_{\Gamma_1} |u|^{m(x)} d\Gamma.$$

Applying (15) and (23), and then using (13) with  $s = m^- + \delta p^-(m^+ - 1) \leq p^-$  and  $s = m^+ + \delta p^-(m^+ - 1) \leq p^-$  we deduce that

$$\begin{aligned} & k^{1-m^-} G^{\delta(m^+-1)}(t) \int_{\Gamma_1} |u|^{m(x)} d\Gamma \\ & \leq k^{1-m^-} \left( \frac{1}{p^-} \int_{\Gamma_1} |u|^{p(x)} d\Gamma \right)^{\delta(m^+-1)} \int_{\Gamma_1} |u|^{m(x)} d\Gamma \\ & \leq k^{1-m^-} (p^-)^{\delta(1-m^+)} C_7 \left\{ \left( \int_{\Gamma_1} |u|^{p(x)} d\Gamma \right)^{\frac{m^-}{p^-} + \delta(m^+-1)} \right. \\ (30) \quad & \quad \left. + \left( \int_{\Gamma_1} |u|^{p(x)} d\Gamma \right)^{\frac{m^+}{p^-} + \delta(m^+-1)} \right\} \\ & = k^{1-m^-} (p^-)^{\delta(1-m^+)} C_7 \left\{ \left( \int_{\Gamma_1} |u|^{p(x)} d\Gamma \right)^{\frac{m^- + \delta p^-(m^+-1)}{p^-}} \right. \\ & \quad \left. + \left( \int_{\Gamma_1} |u|^{p(x)} d\Gamma \right)^{\frac{m^+ + \delta p^-(m^+-1)}{p^-}} \right\} \\ & \leq 2k^{1-m^-} (p^-)^{\delta(1-m^+)} C_5 C_7 \left( \|\nabla u\|_2^2 + \int_{\Gamma_1} |u|^{p(x)} d\Gamma \right). \end{aligned}$$

We note that from the definition of  $E(t)$  and Lemma 3.2,

$$\begin{aligned} \frac{1}{p^-} \int_{\Gamma_1} |u|^{p(x)} d\Gamma & \geq \int_{\Gamma_1} \frac{|u|^{p(x)}}{p(x)} d\Gamma \geq \frac{1}{2} \mu_0 \|\nabla u\|_2^2 - E(t) \\ & \geq \frac{1}{2} \mu_0 \eta_*^2 - d > \frac{1}{2} \mu_0 \eta_1^2 - d = d \left( \frac{2}{p^- - 2} \right) \end{aligned}$$

by the definition of  $d$ . Hence, from the above inequality and the definition of  $E(t)$ , we have

$$\begin{aligned}
 \mu_0 \|\nabla u\|_2^2 &\leq \mu(t) \|\nabla u\|_2^2 \\
 &= 2E(t) - \|u_t\|_2^2 - 2 \int_{\Omega} H(u) dx + 2 \int_{\Gamma_1} \frac{|u|^{p(x)}}{p(x)} d\Gamma \\
 (31) \quad &= 2E_1 - 2G(t) - \|u_t\|_2^2 - 2 \int_{\Omega} H(u) dx + 2 \int_{\Gamma_1} \frac{|u|^{p(x)}}{p(x)} d\Gamma \\
 &< 2d + \frac{2}{p^-} \int_{\Gamma_1} |u|^{p(x)} d\Gamma \\
 &= \int_{\Gamma_1} |u|^{p(x)} d\Gamma.
 \end{aligned}$$

Combining (30) and (31), we obtain

$$\begin{aligned}
 (32) \quad &k^{1-m^-} G^{\delta(m^+-1)}(t) \int_{\Gamma_1} |u|^{m(x)} d\Gamma \\
 &\leq 2k^{1-m^-} (p^-)^{\delta(1-m^+)} C_5 C_7 (\mu_0^{-1} + 1) \int_{\Gamma_1} |u|^{p(x)} d\Gamma.
 \end{aligned}$$

Therefore (29) and (32) yield

$$\begin{aligned}
 (33) \quad &\int_{\Gamma_1} |u_t|^{m(x)-1} |u| d\Gamma \\
 &\leq k G^{-\delta}(t) G'(t) + 2k^{1-m^-} (p^-)^{\delta(1-m^+)} C_5 C_7 (\mu_0^{-1} + 1) \int_{\Gamma_1} |u|^{p(x)} d\Gamma.
 \end{aligned}$$

Combining (26), (27), (28) and (33), we have for  $0 < \epsilon < h_\epsilon$ ,

$$\begin{aligned}
 L'(t) &\geq (1 - \delta - \epsilon k) G^{-\delta}(t) G'(t) + \epsilon \left(1 + \frac{\theta}{2}\right) \|u_t\|_2^2 + \epsilon \theta G(t) \\
 &\quad + \epsilon \left(1 - \frac{\theta}{p^-} - 2k^{1-m^-} (p^-)^{\delta(1-m^+)} C_5 C_7 (\mu_0^{-1} + 1)\right) \int_{\Gamma_1} |u|^{p(x)} d\Gamma.
 \end{aligned}$$

We now choose  $k$  large enough such that

$$1 - \frac{\theta}{p^-} - 2k^{1-m^-} (p^-)^{\delta(1-m^+)} C_5 C_7 (\mu_0^{-1} + 1) \geq 0.$$

Once  $k$  is fixed, we take  $\epsilon$  small enough such that  $1 - \delta - \epsilon k \geq 0$  and  $L(0) = G^{1-\delta}(0) + \epsilon N(0) > 0$ , consequently, we conclude that from (14)

$$L'(t) \geq C_8 (\|u_t\|_2^2 + \|u\|_{p^-}^{p^-} + G(t)),$$

where  $C_8$  is a positive constant, which implies that  $L(t)$  is a positive increasing function. By the same arguments as in [16], page 3036, we have

$$L'(t) \geq C_9 L^{\frac{1}{1-\delta}}(t) \quad \text{for all } t \in (0, T_{\max}),$$

where  $C_9$  is a positive constant. Hence we conclude that  $L(t)$  blows up in finite time and  $u$  also blows up in finite time. Thus the proof of Theorem 2.7 is completed.

## References

- [1] R. Aboulaich, D. Meskine, and A. Souissi, *New diffusion models in image processing*, Comput. Math. Appl. **56** (2008), no. 4, 874–882. <https://doi.org/10.1016/j.camwa.2008.01.017>
- [2] S. Antontsev, *Wave equation with  $p(x,t)$ -Laplacian and damping term: existence and blow-up*, Differ. Equ. Appl. **3** (2011), no. 4, 503–525. <https://doi.org/10.7153/dea-03-32>
- [3] S. Antontsev and S. Shmarev, *Evolution PDEs with nonstandard growth conditions*, Atlantis Studies in Differential Equations, 4, Atlantis Press, Paris, 2015. <https://doi.org/10.2991/978-94-6239-112-3>
- [4] L. Diening, P. Harjulehto, P. Hästö, and M. Růžička, *Lebesgue and Sobolev spaces with variable exponents*, Lecture Notes in Mathematics, 2017, Springer, Heidelberg, 2011. <https://doi.org/10.1007/978-3-642-18363-8>
- [5] X. Fan, *Boundary trace embedding theorems for variable exponent Sobolev spaces*, J. Math. Anal. Appl. **339** (2008), no. 2, 1395–1412. <https://doi.org/10.1016/j.jmaa.2007.08.003>
- [6] V. Georgiev and G. Todorova, *Existence of a solution of the wave equation with nonlinear damping and source terms*, J. Differential Equations **109** (1994), no. 2, 295–308. <https://doi.org/10.1006/jdeq.1994.1051>
- [7] S. Ghegal, I. Hamchi, and S. A. Messaoudi, *Global existence and stability of a nonlinear wave equation with variable-exponent nonlinearities*, Appl. Anal. **99** (2020), no. 8, 1333–1343. <https://doi.org/10.1080/00036811.2018.1530760>
- [8] T. G. Ha, *Blow-up for semilinear wave equation with boundary damping and source terms*, J. Math. Anal. Appl. **390** (2012), no. 1, 328–334. <https://doi.org/10.1016/j.jmaa.2012.01.037>
- [9] T. G. Ha, *Blow-up for wave equation with weak boundary damping and source terms*, Appl. Math. Lett. **49** (2015), 166–172. <https://doi.org/10.1016/j.aml.2015.05.003>
- [10] H. A. Levine, *Instability and nonexistence of global solutions to nonlinear wave equations of the form  $Pu_{tt} = -Au + F(u)$* , Trans. Amer. Math. Soc. **192** (1974), 1–21. <https://doi.org/10.2307/1996814>
- [11] H. A. Levine, *Some additional remarks on the nonexistence of global solutions to nonlinear wave equations*, SIAM J. Math. Anal. **5** (1974), 138–146. <https://doi.org/10.1137/0505015>
- [12] H. A. Levine and J. Serrin, *Global nonexistence theorems for quasilinear evolution equations with dissipation*, Arch. Rational Mech. Anal. **137** (1997), no. 4, 341–361. <https://doi.org/10.1007/s002050050032>
- [13] H. A. Levine and G. Todorova, *Blow up of solutions of the Cauchy problem for a wave equation with nonlinear damping and source terms and positive initial energy*, Proc. Amer. Math. Soc. **129** (2001), no. 3, 793–805. <https://doi.org/10.1090/S0002-9939-00-05743-9>
- [14] S. A. Messaoudi, J. H. Al-Smail, and A. A. Talahmeh, *Decay for solutions of a nonlinear damped wave equation with variable-exponent nonlinearities*, Comput. Math. Appl. **76** (2018), no. 8, 1863–1875. <https://doi.org/10.1016/j.camwa.2018.07.035>
- [15] S. A. Messaoudi and A. A. Talahmeh, *Blowup in solutions of a quasilinear wave equation with variable-exponent nonlinearities*, Math. Methods Appl. Sci. **40** (2017), no. 18, 6976–6986. <https://doi.org/10.1002/mma.4505>

- [16] S. A. Messaoudi, A. A. Talahmeh, and J. H. Al-Smail, *Nonlinear damped wave equation: existence and blow-up*, Comput. Math. Appl. **74** (2017), no. 12, 3024–3041. <https://doi.org/10.1016/j.camwa.2017.07.048>
- [17] L. E. Payne and D. H. Sattinger, *Saddle points and instability of nonlinear hyperbolic equations*, Israel J. Math. **22** (1975), no. 3-4, 273–303. <https://doi.org/10.1007/BF02761595>
- [18] P. Pucci and J. Serrin, *Global nonexistence for abstract evolution equations with positive initial energy*, J. Differential Equations **150** (1998), no. 1, 203–214. <https://doi.org/10.1006/jdeq.1998.3477>
- [19] A. Rahmoune, *Existence and asymptotic stability for the semilinear wave equation with variable-exponent nonlinearities*, J. Math. Phys. **60** (2019), no. 12, 122701, 23 pp. <https://doi.org/10.1063/1.5089879>
- [20] M. Růžička, *Electrorheological fluids: modeling and mathematical theory*, Lecture Notes in Mathematics, 1748, Springer-Verlag, Berlin, 2000. <https://doi.org/10.1007/BFb0104029>
- [21] G. Todorova and E. Vitillaro, *Blow-up for nonlinear dissipative wave equations in  $\mathbb{R}^n$* , J. Math. Anal. Appl. **303** (2005), no. 1, 242–257. <https://doi.org/10.1016/j.jmaa.2004.08.039>
- [22] E. Vitillaro, *A potential well theory for the wave equation with nonlinear source and boundary damping terms*, Glasg. Math. J. **44** (2002), no. 3, 375–395. <https://doi.org/10.1017/S0017089502030045>
- [23] E. Vitillaro, *Global nonexistence theorems for a class of evolution equations with dissipation*, Arch. Ration. Mech. Anal. **149** (1999), no. 2, 155–182. <https://doi.org/10.1007/s002050050171>

TAE GAB HA

DEPARTMENT OF MATHEMATICS AND INSTITUTE OF PURE AND APPLIED MATHEMATICS  
JEONBUK NATIONAL UNIVERSITY  
JEONJU 54896, KOREA  
Email address: [tgha@jbnu.ac.kr](mailto:tgha@jbnu.ac.kr)

SUN-HYE PARK

OFFICE FOR EDUCATION ACCREDITATION  
PUSAN NATIONAL UNIVERSITY  
BUSAN 46241, KOREA  
Email address: [sh-park@pusan.ac.kr](mailto:sh-park@pusan.ac.kr)